

**Updated  
Confidence  
Intervals**

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results – *Bulletin 2*



## Prevalence Table 1 – Northern Ireland

All HSSB Areas

Table 1.1: Northern Ireland – Lifetime Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	20.6	26.7	13.5
	CI+	21.8	29.6	15.4
	CI-	18.3	23.9	11.7
Cannabis	%	16.8	23.3	10.5
	CI+	18.5	26.0	12.1
	CI-	15.3	20.6	9.0
Heroin	%	0.2	0.4	0.0
	CI+	0.4	0.9	0.2
	CI-	0.1	0.1	0.0
Methadone	%	0.1	0.2	0.1
	CI+	0.3	0.6	0.4
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	18.0	16.4	19.5
	CI+	20.2	19.2	22.1
	CI-	15.9	13.9	17.1
Cocaine (total including crack)	%	1.6	2.8	0.5
	CI+	2.2	3.9	1.0
	CI-	1.2	1.9	0.3
Crack	%	0.2	0.4	0.1
	CI+	0.5	1.0	0.3
	CI-	0.1	0.2	0.0
Cocaine powder	%	1.6	2.7	0.5
	CI+	2.1	3.8	0.9
	CI-	1.2	1.8	0.2
Amphetamines	%	4.0	6.3	1.8
	CI+	4.9	7.8	2.6
	CI-	3.3	5.0	1.2
Ecstasy	%	5.8	8.3	3.3
	CI+	6.7	10.0	4.3
	CI-	4.9	6.8	2.4

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	4.5	7.0	2.0
	CI+	5.4	8.7	2.8
	CI-	3.7	5.6	1.4
Magic mushrooms	%	4.3	6.3	2.4
	CI+	5.2	7.9	3.2
	CI-	3.6	4.9	1.8
Solvents	%	2.9	4.1	1.7
	CI+	3.6	5.5	2.4
	CI-	2.3	3.0	1.2
Poppers <sup>3</sup>	%	5.5	7.3	3.8
	CI+	6.6	9.0	5.0
	CI-	4.5	5.9	2.7
Anabolic steroids	%	0.7	0.7	0.8
	CI+	1.1	1.3	1.3
	CI-	0.5	0.3	0.4
Sedatives, tranquillisers, anti-depressants	%	22.1	15.5	28.5
	CI+	23.8	17.7	30.7
	CI-	20.4	13.4	26.5
Alcohol	%	88.2	90.3	86.0
	CI+	89.4	91.9	87.6
	CI-	86.9	88.6	84.3
Tobacco	%	58.6	61.4	55.8
	CI+	60.9	64.6	58.5
	CI-	56.2	58.0	53.1

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

All HSSB Areas

Table 1.2: Northern Ireland – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	6.4	9.7	3.1
	CI+	7.4	11.6	4.1
	CI-	5.4	8.0	2.3
Cannabis	%	5.4	8.7	2.2
	CI+	6.4	10.5	3.1
	CI-	4.5	7.0	1.5
Heroin	%	0.0	0.0	0.0
	CI+	0.1	0.2	0.2
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.1
	CI+	0.2	0.2	0.3
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	8.0	6.9	9.0
	CI+	9.3	8.5	10.6
	CI-	6.8	5.5	7.6
Cocaine (total including crack)	%	0.5	1.0	0.1
	CI+	0.9	1.8	0.4
	CI-	0.3	0.4	0.0
Crack	%	0.0	0.1	0.0
	CI+	0.2	0.5	0.2
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.5	0.9	0.1
	CI+	0.8	1.6	0.4
	CI-	0.2	0.4	0.0
Amphetamines	%	0.8	1.4	0.2
	CI+	1.3	2.3	0.5
	CI-	0.5	0.8	0.0
Ecstasy	%	1.6	2.7	0.6
	CI+	2.2	3.9	1.1
	CI-	1.2	1.8	0.3

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.1	0.0	0.1
	CI+	0.2	0.2	0.5
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.2	0.4	0.3
	CI+	0.5	0.9	0.9
	CI-	0.1	0.1	0.1
Solvents	%	0.2	0.3	0.3
	CI+	0.4	0.8	0.9
	CI-	0.0	0.0	0.1
Poppers <sup>3</sup>	%	0.5	0.7	1.1
	CI+	0.8	1.4	1.9
	CI-	0.2	0.3	0.5
Anabolic steroids	%	0.3	0.3	0.4
	CI+	0.6	0.8	1.0
	CI-	0.2	0.1	0.1
Sedatives, tranquillisers, anti-depressants	%	12.6	8.7	16.5
	CI+	13.9	10.3	18.3
	CI-	11.4	7.3	14.8
Alcohol	%	79.3	82.2	83.4
	CI+	80.7	84.1	85.4
	CI-	77.8	80.1	81.1
Tobacco	%	40.2	40.9	43.2
	CI+	42.6	44.0	46.3
	CI-	37.9	37.7	40.1

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

All HSSB Areas

Table 1.3: Northern Ireland – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	3.4	5.7	1.1
	CI+	4.2	7.3	1.7
	CI-	2.6	4.3	0.6
Cannabis	%	2.9	5.1	0.8
	CI+	3.7	6.7	1.3
	CI-	2.3	3.8	0.4
Heroin	%	0.0	0.0	0.0
	CI+	0.1	0.2	0.2
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.1
	CI+	0.2	0.2	0.3
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	4.1	3.1	5.2
	CI+	4.9	4.0	6.3
	CI-	3.4	2.3	4.2
Cocaine (total including crack)	%	0.1	0.2	0.0
	CI+	0.3	0.6	0.2
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.1	0.2	0.2
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.1	0.2	0.0
	CI+	0.3	0.6	0.2
	CI-	0.0	0.0	0.0
Amphetamines	%	0.2	0.5	0.0
	CI+	0.5	1.0	0.2
	CI-	0.1	0.2	0.0
Ecstasy	%	0.5	0.9	0.1
	CI+	0.9	1.6	0.5
	CI-	0.3	0.4	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.1	0.2	0.3
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.1	0.0
	CI+	0.2	0.4	0.2
	CI-	0.0	0.0	0.0
Solvents	%	0.1	0.1	0.1
	CI+	0.2	0.3	0.3
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.2	0.2	0.3
	CI+	0.4	0.7	0.9
	CI-	0.0	0.0	0.1
Anabolic steroids	%	0.1	0.1	0.1
	CI+	0.3	0.3	0.5
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	9.8	7.0	12.5
	CI+	10.9	8.5	14.1
	CI-	8.7	5.7	11.1
Alcohol	%	67.6	72.7	62.5
	CI+	69.3	75.0	64.7
	CI-	65.8	70.3	60.3
Tobacco	%	37.0	37.0	37.1
	CI+	39.4	40.0	39.8
	CI-	34.7	33.9	34.4

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 2 – Northern Ireland

Eastern Health and Social Services Board (EHSSB)

Table 2.1: EHSSB – Lifetime Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64		All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males				15-64	Males		
Any illegal drugs <sup>1</sup>	%	22.5	29.8	15.5	33.7	13.7	LSD	%	5.0
	CI+	25.8	35.2	19.0	39.6	16.6	CI+	6.5	8.1
	CI-	19.4	24.8	12.5	28.2	11.2	CI-	3.8	2.2
Cannabis	%	18.9	26.2	12.0	27.4	12.3	Magic mushrooms	%	4.3
	CI+	22.0	31.5	14.9	32.9	15.1	CI+	5.5	5.7
	CI-	16.1	21.3	9.5	22.3	9.8	CI-	3.2	3.7
Heroin	%	0.1	0.2	0.0	0.2	0.0	Solvents	%	2.4
	CI+	0.4	0.9	0.5	1.0	0.4	CI+	3.6	3.2
	CI-	0.0	0.0	0.0	0.0	0.0	CI-	1.6	2.0
Methadone	%	0.2	0.3	0.1	0.3	0.1	Poppers <sup>3</sup>	%	6.3
	CI+	0.7	1.7	0.7	1.9	0.6	CI+	8.5	8.2
	CI-	0.0	0.0	0.0	0.0	0.0	CI-	4.5	4.7
Other opiates <sup>2</sup>	%	22.9	22.6	23.2	22.5	23.3	Anabolic steroids	%	0.8
	CI+	27.5	28.5	28.1	28.6	28.0	CI+	1.5	1.2
	CI-	18.9	17.5	18.9	17.2	19.0	CI-	0.4	0.7
Cocaine (total including crack)	%	2.4	4.0	0.9	4.5	0.8	Sedatives, tranquillisers, anti-depressants	%	22.3
	CI+	3.5	6.5	1.8	6.9	1.7	CI+	25.3	29.3
	CI-	1.6	2.3	0.3	2.7	0.3	CI-	19.6	16.4
Crack	%	0.1	0.0	0.1	0.0	0.1	Alcohol	%	90.9
	CI+	0.4	0.6	0.7	0.7	0.6	CI+	92.7	90.1
	CI-	0.0	0.0	0.0	0.0	0.0	CI-	92.4	91.6
Cocaine powder	%	2.3	4.0	0.7	4.5	0.7	Tobacco	%	59.5
	CI+	3.5	6.5	1.6	6.9	1.6	CI+	61.7	92.9
	CI-	1.5	2.3	0.3	2.7	0.2	CI-	66.5	93.4
Amphetamines	%	4.9	7.4	2.6	9.4	1.4	CI-	89.3	86.5
	CI+	6.5	10.2	4.1	12.7	2.4	CI-	55.9	89.5
	CI-	3.6	5.1	1.5	6.8	0.7	CI-	56.7	61.5
Ecstasy	%	7.1	10.1	4.2	13.9	1.7	CI-	52.6	65.9
	CI+	8.8	13.3	6.0	17.6	2.9	CI-	51.4	66.5
	CI-	5.6	7.4	2.8	10.8	0.9	CI-	57.0	65.9

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Eastern Health and Social Services Board (EHSSB)**

Table 2.2: EHSSB – Last Year Prevalence (%)

	All Adults 15-64			Young Adults 15-34	Older Adults 35-64
		Males	Females		
Any illegal drugs <sup>1</sup>	%	7.3	11.0	3.9	13.3
	CI+	9.2	14.4	6.0	17.2
	CI-	5.7	8.1	2.3	10.0
Cannabis	%	6.0	9.3	2.9	10.8
	CI+	7.9	12.6	5.0	14.8
	CI-	4.4	6.6	1.5	7.6
Heroin	%	0.0	0.0	0.0	0.0
	CI+	0.3	0.6	0.5	0.7
	CI-	0.0	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0	0.0
	CI+	0.3	0.6	0.5	0.7
	CI-	0.0	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	9.4	8.5	10.2	8.7
	CI+	11.7	11.7	13.1	12.2
	CI-	7.4	5.9	7.7	6.0
Cocaine (total including crack)	%	1.0	1.9	0.2	2.0
	CI+	2.0	4.1	1.0	4.0
	CI-	0.5	0.7	0.0	0.8
Crack	%	0.0	0.0	0.0	0.0
	CI+	0.3	0.6	0.5	0.7
	CI-	0.0	0.0	0.0	0.0
Cocaine powder	%	1.0	1.9	0.2	2.0
	CI+	2.0	4.1	1.0	4.0
	CI-	0.5	0.7	0.0	0.8
Amphetamines	%	1.5	2.7	0.3	2.8
	CI+	2.6	4.9	1.0	5.2
	CI-	0.8	1.3	0.0	1.3
Ecstasy	%	2.3	3.6	1.2	4.6
	CI+	3.4	5.8	2.3	6.9
	CI-	1.5	2.0	0.5	2.9

	All Adults 15-64			Young Adults 15-34	Older Adults 35-64
		Males	Females		
LSD	%	0.1	0.0	0.1	0.0
	CI+	0.4	0.6	0.7	0.9
	CI-	0.0	0.0	0.0	0.0
Magic mushrooms	%	0.2	0.3	0.0	0.0
	CI+	0.6	1.3	0.5	0.7
	CI-	0.0	0.0	0.0	0.0
Solvents	%	0.3	0.6	0.0	0.6
	CI+	1.0	2.1	0.5	2.3
	CI-	0.0	0.1	0.0	0.1
Poppers <sup>3</sup>	%	0.6	0.8	0.3	1.3
	CI+	1.3	2.5	1.0	3.1
	CI-	0.2	0.2	0.0	0.4
Anabolic steroids	%	0.3	0.4	0.3	0.3
	CI+	0.9	1.8	0.9	1.9
	CI-	0.1	0.0	0.0	0.1
Sedatives, tranquillisers, anti-depressants	%	13.3	8.9	17.3	9.2
	CI+	15.4	11.8	20.2	12.1
	CI-	11.3	6.6	14.8	6.8
Alcohol	%	82.3	85.7	79.2	84.3
	CI+	84.3	88.3	81.9	87.7
	CI-	80.2	82.6	76.1	80.5
Tobacco	%	40.3	40.2	40.4	42.5
	CI+	44.3	45.3	45.0	48.1
	CI-	36.4	35.2	36.0	37.1

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Eastern Health and Social Services Board (EHSSB)**

Table 2.3: EHSSB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	3.8	6.6	1.2
	CI+	5.3	9.6	2.5
	CI-	2.6	4.3	0.5
Cannabis	%	3.4	5.9	1.0
	CI+	4.9	8.9	2.4
	CI-	2.2	3.7	0.3
Heroin	%	0.0	0.0	0.0
	CI+	0.3	0.6	0.5
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.3	0.6	0.5
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	4.6	4.0	5.1
	CI+	5.9	5.9	7.0
	CI-	3.5	2.6	3.6
Cocaine (total including crack)	%	0.1	0.3	0.0
	CI+	0.8	1.7	0.5
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.3	0.6	0.5
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.1	0.3	0.0
	CI+	0.8	1.7	0.5
	CI-	0.0	0.0	0.0
Amphetamines	%	0.4	0.8	0.0
	CI+	0.9	2.0	0.5
	CI-	0.1	0.2	0.0
Ecstasy	%	0.5	0.9	0.2
	CI+	1.2	2.4	1.0
	CI-	0.2	0.2	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.1	0.0	0.1
	CI+	0.4	0.6	0.7
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.1	0.2	0.0
	CI+	0.5	1.0	0.5
	CI-	0.0	0.0	0.0
Solvents	%	0.0	7.3	0.0
	CI+	0.3	10.0	0.5
	CI-	0.0	5.1	0.0
Poppers <sup>3</sup>	%	0.2	0.3	0.4
	CI+	0.8	1.7	1.8
	CI-	0.0	0.0	0.0
Anabolic steroids	%	0.1	0.0	0.2
	CI+	0.5	0.6	0.9
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	10.1	0.0	12.7
	CI+	12.0	0.6	15.3
	CI-	8.3	0.0	10.5
Alcohol	%	71.9	78.6	65.5
	CI+	74.4	81.8	68.8
	CI-	69.2	75.1	62.1
Tobacco	%	37.4	36.5	38.3
	CI+	41.5	41.8	43.0
	CI-	33.4	31.4	33.7

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 3 – Northern Ireland

Northern Health and Social Services Board (NHSSB)

Table 3.1: NHSSB – Lifetime Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males			15-64	Males		
Any illegal drugs <sup>1</sup>	%	17.8	22.9	12.7	29.3	9.1	LSD	%
	CI+	21.5	28.6	16.9	36.4	12.1		CI+
	CI-	14.5	17.9	9.2	22.9	6.6		CI-
Cannabis	%	14.4	19.5	9.3	22.8	8.1	Magic mushrooms	%
	CI+	17.4	24.7	12.7	28.6	11.1		CI+
	CI-	11.7	15.0	6.6	17.6	5.7		CI-
Heroin	%	0.4	0.8	0.0	0.4	0.4	Solvents	%
	CI+	1.2	2.5	0.8	2.3	1.5		CI+
	CI-	0.1	0.2	0.0	0.0	0.1		CI-
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%
	CI+	0.4	0.9	0.8	1.1	0.7		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Other opiates <sup>2</sup>	%	18.0	15.3	20.7	20.8	15.9	Anabolic steroids	%
	CI+	22.4	20.8	26.2	27.1	20.7		CI+
	CI-	14.1	10.7	15.9	15.4	11.8		CI-
Cocaine (total including crack)	%	1.0	2.1	0.0	1.3	0.9	Sedatives, tranquillisers, anti-depressants	%
	CI+	2.2	4.4	0.8	3.5	2.1		CI+
	CI-	0.4	0.8	0.0	0.3	0.3		CI-
Crack	%	0.3	0.6	0.0	0.4	0.2	Alcohol	%
	CI+	1.0	2.2	0.8	2.3	1.2		CI+
	CI-	0.0	0.1	0.0	0.0	0.0		CI-
Cocaine powder	%	1.0	2.1	0.0	1.3	0.9	Tobacco	%
	CI+	2.2	4.4	0.8	3.5	2.1		CI+
	CI-	0.4	0.8	0.0	0.3	0.3		CI-
Amphetamines	%	4.1	6.3	1.8	6.4	2.3		
	CI+	5.7	9.2	4.2	10.1	4.0		
	CI-	2.8	4.1	0.6	3.7	1.2		
Ecstasy	%	5.9	8.7	3.2	11.3	1.9		
	CI+	8.1	12.3	6.3	16.3	3.5		
	CI-	4.2	5.9	1.4	7.4	0.9		

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

Northern Health and Social Services Board (NHSSB)

Table 3.2: NHSSB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males			15-64	Males		
Any illegal drugs <sup>1</sup>	%	6.2	9.8	2.7	11.7	2.1	LSD	%
	CI+	8.6	14.0	4.7	17.0	3.8		CI+
	CI-	4.3	6.5	1.4	7.5	1.0		CI-
Cannabis	%	5.4	9.1	1.7	9.8	2.1	Magic mushrooms	%
	CI+	7.5	13.2	3.3	14.4	3.8		CI+
	CI-	3.8	6.0	0.7	6.2	1.0		CI-
Heroin	%	0.0	0.0	0.0	0.0	0.0	Solvents	%
	CI+	0.4	0.9	0.8	1.1	0.7		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%
	CI+	0.4	0.9	0.8	1.1	0.7		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Other opiates <sup>2</sup>	%	9.5	8.4	10.6	10.2	9.0	Anabolic steroids	%
	CI+	12.5	12.0	14.1	14.1	12.4		CI+
	CI-	7.0	5.6	7.6	7.0	6.3		CI-
Cocaine (total including crack)	%	0.0	0.0	0.0	0.0	0.0	Sedatives, tranquillisers, anti-depressants	%
	CI+	0.4	0.9	0.8	1.1	0.7		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Crack	%	0.0	0.0	0.0	0.0	0.0	Alcohol	%
	CI+	0.4	0.9	0.8	1.1	0.7		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Cocaine powder	%	0.0	0.0	0.0	0.0	0.0	Tobacco	%
	CI+	0.4	0.9	0.8	1.1	0.7		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Amphetamines	%	0.6	0.9	0.2	1.0	0.2		
	CI+	1.4	2.7	1.2	3.2	1.2		
	CI-	0.2	0.2	0.0	0.2	0.0		
Ecstasy	%	1.5	3.1	0.0	3.0	0.4		
	CI+	3.0	6.0	0.8	6.2	1.5		
	CI-	0.7	1.3	0.0	1.1	0.1		

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

Northern Health and Social Services Board (NHSSB)

Table 3.3: NHSSB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	3.8	6.5	1.0
	CI+	5.9	10.4	2.6
	CI-	2.2	3.7	0.3
Cannabis	%	3.5	6.2	0.8
	CI+	5.3	9.9	2.2
	CI-	2.1	3.5	0.2
Heroin	%	0.0	0.0	0.0
	CI+	0.4	0.9	0.8
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.4	0.9	0.8
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	5.2	3.8	6.6
	CI+	7.2	6.3	9.7
	CI-	3.6	2.1	4.3
Cocaine (total including crack)	%	0.0	0.0	0.0
	CI+	0.4	0.9	0.8
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.4	0.9	0.8
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.0	0.0	0.0
	CI+	0.4	0.9	0.8
	CI-	0.0	0.0	0.0
Amphetamines	%	0.2	0.3	0.0
	CI+	0.9	1.9	0.8
	CI-	0.0	0.0	0.0
Ecstasy	%	1.1	2.2	0.0
	CI+	2.1	4.3	0.8
	CI-	0.5	0.9	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.4	0.9	1.1
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.4	0.9	1.1
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.4	0.9	1.1
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.1	0.3	0.3
	CI+	0.8	1.5	2.0
	CI-	0.0	0.0	0.0
Anabolic steroids	%	0.0	0.0	0.0
	CI+	0.4	0.9	1.1
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	10.5	13.5	5.6
	CI+	13.0	10.7	9.5
	CI-	8.3	4.9	3.0
Alcohol	%	64.5	61.3	65.0
	CI+	67.9	65.7	71.4
	CI-	61.0	56.8	58.1
Tobacco	%	35.3	35.3	37.5
	CI+	40.2	40.5	44.1
	CI-	30.6	29.2	31.2

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 4 – Northern Ireland

Southern Health and Social Services Board (SHSSB)

Table 4.1: SHSSB – Lifetime Prevalence (%)

	All Adults 15-64		Young Adults 15-34	Older Adults 35-64		All Adults 15-64		Young Adults 15-34	Older Adults 35-64
	Males	Females				Males	Females		
Any illegal drugs <sup>1</sup>	%	19.1	26.4	11.8	28.8	11.4			
	CI+	22.8	32.5	16.3	35.4	15.0			
	CI-	15.8	20.9	8.1	22.9	8.4			
Cannabis	%	16.3	23.0	9.5	23.9	10.2			
	CI+	19.6	28.9	13.1	30.1	13.6			
	CI-	13.3	17.8	6.6	18.5	7.3			
Heroin	%	0.0	0.0	0.0	0.0	0.0			
	CI+	0.6	1.2	1.1	1.4	0.9			
	CI-	0.0	0.0	0.0	0.0	0.0			
Methadone	%	0.2	0.0	0.3	0.0	0.3			
	CI+	0.9	1.2	1.6	1.4	1.5			
	CI-	0.0	0.0	0.0	0.0	0.0			
Other opiates <sup>2</sup>	%	13.1	11.2	15.0	9.9	15.6			
	CI+	16.3	15.3	19.6	14.1	19.6			
	CI-	10.3	7.9	11.1	6.6	12.1			
Cocaine (total including crack)	%	0.8	1.4	0.2	1.5	0.2			
	CI+	2.0	3.5	1.5	3.8	1.4			
	CI-	0.2	0.4	0.0	0.4	0.0			
Crack	%	0.0	0.0	0.0	0.0	0.0			
	CI+	0.6	1.2	1.1	1.4	0.9			
	CI-	0.0	0.0	0.0	0.0	0.0			
Cocaine powder	%	0.8	1.4	0.2	1.5	0.2			
	CI+	2.0	3.5	1.5	3.8	1.4			
	CI-	0.2	0.4	0.0	0.4	0.0			
Amphetamines	%	2.3	4.1	0.5	3.4	1.4			
	CI+	3.9	7.4	1.9	6.4	3.2			
	CI-	1.2	2.0	0.0	1.6	0.5			
Ecstasy	%	3.2	4.9	1.5	6.1	0.9			
	CI+	5.2	8.5	3.8	10.2	2.5			
	CI-	1.8	2.5	0.4	3.2	0.2			
LSD	%		3.4		4.4	2.4	6.5	0.9	
	CI+		5.6		8.7	4.8	11.0	2.7	
	CI-		1.9		1.8	1.0	3.4	0.2	
Magic mushrooms	%		2.9		4.1	1.8	4.3	1.9	
	CI+		4.5		7.0	3.8	7.8	3.8	
	CI-		1.8		2.1	0.7	2.1	0.8	
Solvents	%		3.2		3.7	2.7	6.1	0.9	
	CI+		5.1		6.9	5.0	10.2	2.5	
	CI-		1.9		1.7	1.3	3.3	0.2	
Poppers <sup>3</sup>	%		3.4		5.1	1.8	7.1	0.5	
	CI+		5.4		8.6	3.7	11.3	1.8	
	CI-		2.0		2.7	0.7	4.2	0.0	
Anabolic steroids	%		0.4		0.3	0.5	0.3	0.5	
	CI+		1.3		2.0	2.0	1.9	1.9	
	CI-		0.1		0.0	0.1	0.0	0.1	
Sedatives, tranquillisers, anti-depressants	%		21.2		16.4	26.1	17.2	24.5	
	CI+		25.9		22.3	31.8	23.5	29.6	
	CI-		17.1		11.5	21.0	12.0	19.9	
Alcohol	%		85.7		88.9	82.5	89.3	82.9	
	CI+		88.7		92.7	86.4	92.9	86.8	
	CI-		82.4		84.1	78.0	84.6	78.5	
Tobacco	%		57.3		61.8	52.8	54.4	59.7	
	CI+		63.1		69.6	58.8	61.1	67.0	
	CI-		51.4		53.5	46.7	47.6	52.0	

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Southern Health and Social Services Board (SHSSB)**

Table 4.2: SHSSB – Last Year Prevalence (%)

	All Adults 15-64		Males	Females	Young Adults 15-34	Older Adults 35-64
	15-64	35-64			15-34	35-64
Any illegal drugs <sup>1</sup>	%	5.1	8.2	1.8	9.6	1.5
	CI+	7.3	12.8	3.9	14.3	3.5
	CI-	3.3	4.9	0.7	6.0	0.5
Cannabis	%	4.3	7.9	0.7	7.9	1.5
	CI+	6.6	12.5	2.3	12.9	3.5
	CI-	2.6	4.6	0.1	4.3	0.5
Heroin	%	0.0	0.0	0.0	0.0	0.0
	CI+	0.6	1.2	1.1	1.4	0.9
	CI-	0.0	0.0	0.0	0.0	0.0
Methadone	%	0.2	0.0	0.3	0.0	0.3
	CI+	0.9	1.2	1.6	1.4	1.5
	CI-	0.0	0.0	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	5.5	4.1	6.9	4.0	6.7
	CI+	8.2	7.1	10.5	7.9	9.7
	CI-	3.5	2.1	4.2	1.6	4.4
Cocaine (total including crack)	%	0.2	0.3	0.0	0.4	0.0
	CI+	0.9	1.9	1.1	2.1	0.9
	CI-	0.0	0.0	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0	0.0	0.0
	CI+	0.6	1.2	1.1	1.4	0.9
	CI-	0.0	0.0	0.0	0.0	0.0
Cocaine powder	%	0.2	0.3	0.0	0.4	0.0
	CI+	0.9	1.9	1.1	2.1	0.9
	CI-	0.0	0.0	0.0	0.0	0.0
Amphetamines	%	0.0	0.0	0.0	0.0	0.0
	CI+	0.6	1.2	1.1	1.4	0.9
	CI-	0.0	0.0	0.0	0.0	0.0
Ecstasy	%	0.5	0.7	0.4	0.9	0.2
	CI+	1.5	2.6	2.0	3.3	1.4
	CI-	0.1	0.1	0.0	0.1	0.0

	All Adults 15-64		Males	Females	Young Adults 15-34	Older Adults 35-64
	15-64	35-64			15-34	35-64
LSD	%	0.2	0.0	0.4	0.4	0.0
	CI+	1.1	1.2	2.0	2.5	0.9
	CI-	0.0	0.0	0.0	0.0	0.0
Magic mushrooms	%	0.4	0.9	0.0	1.0	0.0
	CI+	1.5	3.1	1.1	3.4	0.9
	CI-	0.1	0.1	0.0	0.1	0.0
Solvents	%	0.2	0.0	0.4	0.4	0.0
	CI+	1.1	1.2	2.0	2.5	0.9
	CI-	0.0	0.0	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.4	0.9	0.0	1.0	0.0
	CI+	1.5	3.1	1.1	3.5	0.9
	CI-	0.1	0.1	0.0	0.1	0.0
Anabolic steroids	%	0.3	0.0	0.5	0.3	0.2
	CI+	1.0	1.2	2.0	1.9	1.4
	CI-	0.0	0.0	0.1	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	11.0	8.9	13.2	6.7	14.5
	CI+	14.1	12.7	17.6	10.6	18.6
	CI-	8.4	6.0	9.6	3.9	11.0
Alcohol	%	76.5	79.9	73.1	85.0	69.7
	CI+	79.7	84.3	78.1	89.4	74.3
	CI-	73.1	74.8	67.6	79.6	64.9
Tobacco	%	38.6	41.9	35.3	41.2	36.5
	CI+	43.9	49.4	41.6	47.3	43.4
	CI-	33.5	34.6	29.3	35.2	30.0

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Southern Health and Social Services Board (SHSSB)**

Table 4.3: SHSSB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	2.1	3.5	0.8
	CI+	3.6	6.4	2.6
	CI-	1.2	1.7	0.1
Cannabis	%	1.5	3.1	0.0
	CI+	2.9	5.8	1.1
	CI-	0.7	1.4	0.0
Heroin	%	0.0	0.0	0.0
	CI+	0.6	1.2	1.1
	CI-	0.0	0.0	0.0
Methadone	%	0.2	0.0	0.3
	CI+	0.9	1.2	1.6
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	2.8	1.2	4.5
	CI+	4.4	3.1	7.2
	CI-	1.7	0.3	2.6
Cocaine (total including crack)	%	0.0	0.0	0.0
	CI+	0.6	1.2	1.1
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.6	1.2	1.1
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.0	0.0	0.0
	CI+	0.6	1.2	1.1
	CI-	0.0	0.0	0.0
Amphetamines	%	0.0	0.0	0.0
	CI+	0.6	1.2	1.1
	CI-	0.0	0.0	0.0
Ecstasy	%	0.2	0.0	0.4
	CI+	1.1	1.2	2.0
	CI-	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.6	1.2	1.1
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.6	1.2	1.1
	CI-	0.0	0.0	0.0
Solvents	%	0.2	0.0	0.4
	CI+	1.1	1.2	2.0
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.2	0.4	0.5
	CI+	1.2	2.4	1.1
	CI-	0.0	0.0	0.0
Anabolic steroids	%	0.0	0.0	0.0
	CI+	0.6	1.2	1.1
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	8.7	6.8	10.7
	CI+	11.4	10.2	14.6
	CI-	6.5	4.2	7.5
Alcohol	%	64.5	70.3	58.5
	CI+	68.4	75.9	64.3
	CI-	60.3	64.3	52.7
Tobacco	%	35.1	37.2	33.0
	CI+	40.3	44.5	39.1
	CI-	30.3	30.3	27.4

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 5 – Northern Ireland

Western Health and Social Services Board (WHSSB)

Table 5.1: WHSSB – Lifetime Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64		All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males				15-64	Males		
Any illegal drugs <sup>1</sup>	%	18.7	25.5	11.8	28.9	9.9	LSD	%	4.1
	CI+	22.6	32.0	15.7	36.3	14.4		CI+	6.2
	CI-	15.2	19.8	8.5	22.3	6.4		CI-	2.1
Cannabis	%	16.3	22.5	9.9	24.5	9.2	Magic mushrooms	%	7.9
	CI+	20.4	29.2	13.9	32.4	13.5		CI+	13.2
	CI-	12.7	16.8	6.8	17.7	5.9		CI-	4.2
Heroin	%	0.4	0.9	0.0	1.0	0.0	Solvents	%	0.9
	CI+	1.6	3.4	1.1	3.5	1.1		CI+	7.5
	CI-	0.1	0.1	0.0	0.1	0.0		CI-	1.8
Methadone	%	0.2	0.4	0.0	0.4	0.0	Poppers <sup>3</sup>	%	8.2
	CI+	1.0	2.1	1.1	2.2	1.1		CI+	12.5
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	3.9
Other opiates <sup>2</sup>	%	11.4	9.6	13.3	11.8	11.1	Anabolic steroids	%	1.6
	CI+	16.4	15.7	19.4	18.4	16.1		CI+	3.3
	CI-	7.5	5.3	8.6	7.0	7.2		CI-	2.1
Cocaine (total including crack)	%	1.6	2.5	0.8	3.3	0.2	Sedatives, tranquillisers, anti-depressants	%	7.9
	CI+	3.0	5.3	2.5	6.3	1.5		CI+	26.6
	CI-	0.8	0.9	0.1	1.5	0.0		CI-	13.3
Crack	%	0.9	1.8	0.0	1.9	0.0		CI-	26.5
	CI+	2.2	4.5	1.1	4.8	1.1		CI-	31.6
	CI-	0.3	0.5	0.0	0.5	0.0		CI-	9.1
Cocaine powder	%	1.5	2.1	0.8	2.9	0.2	Alcohol	%	21.9
	CI+	2.8	4.9	2.5	5.8	1.5		CI+	91.6
	CI-	0.7	0.7	0.1	1.2	0.0		CI-	83.7
Amphetamines	%	3.8	6.1	1.4	7.5	0.6		CI-	83.7
	CI+	6.4	11.0	3.3	13.3	2.1		CI-	94.9
	CI-	2.0	3.0	0.4	3.7	0.1		CI-	87.8
Ecstasy	%	5.2	7.4	3.1	10.9	0.3	Tobacco	%	79.9
	CI+	7.9	12.5	5.6	17.1	1.7		CI+	87.3
	CI-	3.3	3.9	1.5	6.4	0.0		CI-	78.9

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Western Health and Social Services Board (WHSSB)**

Table 5.2: WHSSB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males			15-64	Males		
Any illegal drugs <sup>1</sup>	%	5.8	8.1	3.6	10.2	2.1	LSD	%
	CI+	8.3	13.2	6.2	15.5	4.7	CI+	0.0
	CI-	3.9	4.5	1.8	6.2	0.8	CI-	0.6
Cannabis	%	5.3	7.4	3.2	9.4	1.8	Magic mushrooms	%
	CI+	7.8	12.6	5.8	14.6	4.3	CI+	0.0
	CI-	3.5	3.9	1.6	5.6	0.6	CI-	0.0
Heroin	%	0.0	0.0	0.0	0.0	0.0	Solvents	%
	CI+	0.6	1.4	1.1	1.4	1.1	CI+	0.2
	CI-	0.0	0.0	0.0	0.0	0.0	CI-	0.9
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%
	CI+	0.6	1.4	1.1	1.4	1.1	CI+	0.2
	CI-	0.0	0.0	0.0	0.0	0.0	CI-	1.0
Other opiates <sup>2</sup>	%	4.9	4.0	5.9	5.0	4.9	Anabolic steroids	%
	CI+	8.4	8.2	9.6	9.9	8.3	CI+	0.9
	CI-	2.6	1.6	3.3	2.1	2.6	CI-	2.1
Cocaine (total including crack)	%	0.4	0.9	0.0	1.0	0.0	Sedatives, tranquillisers, anti-depressants	%
	CI+	1.6	3.3	1.1	3.5	1.1	CI+	11.2
	CI-	0.1	0.1	0.0	0.1	0.0	CI-	14.1
Crack	%	0.3	0.5	0.0	0.6	0.0	Alcohol	%
	CI+	1.4	3.1	1.1	3.2	1.1	CI+	78.4
	CI-	0.0	0.0	0.0	0.0	0.0	CI-	81.9
Cocaine powder	%	0.2	0.4	0.0	0.4	0.0	Tobacco	%
	CI+	1.0	2.1	1.1	2.2	1.1	CI+	44.7
	CI-	0.0	0.0	0.0	0.0	0.0	CI-	50.1
Amphetamines	%	0.4	0.9	0.0	1.0	0.0		
	CI+	1.6	3.3	1.1	3.5	1.1		
	CI-	0.1	0.1	0.0	0.1	0.0		
Ecstasy	%	1.4	2.5	0.3	3.0	0.0		
	CI+	3.3	6.4	1.8	7.2	1.1		
	CI-	0.5	0.6	0.0	0.9	0.0		

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Western Health and Social Services Board (WHSSB)**

Table 5.3: WHSSB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males			15-64	Males		
Any illegal drugs <sup>1</sup>	%	3.0	4.7	1.2	4.9	1.3	LSD	%
	CI+	5.1	9.3	3.0	9.1	3.9		CI+
	CI-	1.5	2.0	0.3	2.3	0.2		CI-
Cannabis	%	2.6	4.1	1.2	4.5	1.0	Magic mushrooms	%
	CI+	4.8	8.7	3.0	8.7	3.7		CI+
	CI-	1.3	1.5	0.3	2.0	0.1		CI-
Heroin	%	0.0	0.0	0.0	0.0	0.0	Solvents	%
	CI+	0.6	1.4	1.1	1.4	1.1		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%
	CI+	0.6	1.4	1.1	1.4	1.1		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Other opiates <sup>2</sup>	%	2.8	1.8	3.9	1.8	3.7	Anabolic steroids	%
	CI+	4.9	4.6	6.6	5.4	6.3		CI+
	CI-	1.5	0.5	2.1	0.3	1.9		CI-
Cocaine (total including crack)	%	0.2	0.4	0.0	0.4	0.0	Sedatives, tranquillisers, anti-depressants	%
	CI+	1.0	2.1	1.1	2.2	1.1		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Crack	%	0.0	0.0	0.0	0.0	0.0	Alcohol	%
	CI+	0.6	1.4	1.1	1.4	1.1		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Cocaine powder	%	0.2	0.4	0.0	0.4	0.0	Tobacco	%
	CI+	1.0	2.1	1.1	2.2	1.1		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Amphetamines	%	0.2	0.4	0.0	0.4	0.0		
	CI+	1.0	2.1	1.1	2.2	1.1		
	CI-	0.0	0.0	0.0	0.0	0.0		
Ecstasy	%	0.0	0.0	0.0	0.0	0.0		
	CI+	0.6	1.4	1.1	1.4	1.1		
	CI-	0.0	0.0	0.0	0.0	0.0		

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, morphine, Temgesic®, codeine, Kapake®, DFs, diffs, dikes and peach.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 6 – Ireland

All HB Areas

Table 6.1: Ireland – Lifetime Prevalence (%)

	All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		
Any illegal drugs <sup>1</sup>	%	18.5	24.0	13.1	26.0	11.8	LSD	%	2.9	4.4	1.4	4.6	1.4
	CI+	20.1	26.6	14.6	28.5	13.3		CI+	3.5	5.5	2.0	5.7	1.9
	CI-	17.0	21.5	11.6	23.5	10.5		CI-	2.4	3.4	1.0	3.6	1.0
Cannabis	%	17.4	22.4	12.3	24.0	11.4	Magic mushrooms	%	3.9	5.7	2.0	5.9	2.0
	CI+	18.9	24.9	13.8	26.5	12.8		CI+	4.5	6.9	2.7	7.3	2.6
	CI-	15.9	20.0	10.9	21.6	10.0		CI-	3.3	4.7	1.5	4.7	1.6
Heroin	%	0.5	0.7	0.3	0.7	0.3	Solvents	%	1.7	2.2	1.1	3.3	0.2
	CI+	0.8	1.2	0.6	1.3	0.6		CI+	2.2	3.1	1.7	4.4	0.4
	CI-	0.3	0.3	0.1	0.4	0.1		CI-	1.3	1.5	0.7	2.4	0.1
Methadone	%	0.3	0.4	0.3	0.6	0.1	Poppers <sup>3</sup>	%	2.6	3.9	1.3	4.7	0.7
	CI+	0.6	0.8	0.7	1.1	0.3		CI+	3.2	5.0	1.9	6.1	1.1
	CI-	0.2	0.1	0.1	0.3	0.0		CI-	2.1	2.9	0.9	3.6	0.4
Other opiates <sup>2</sup>	%	3.0	2.0	4.0	3.0	3.1	Sedatives, tranquillisers, anti-depressants	%	12.1	9.3	15.0	8.2	15.7
	CI+	3.7	2.8	5.0	4.1	3.8		CI+	13.2	10.6	16.5	9.6	17.2
	CI-	2.4	1.4	3.1	2.1	2.4		CI-	11.2	8.0	13.7	6.9	14.3
Cocaine (total including crack)	%	3.0	4.3	1.6	4.7	1.4	Alcohol	%	90.2	92.5	87.8	92.1	88.5
	CI+	3.6	5.4	2.3	5.9	2.0		CI+	91.1	93.7	89.0	93.4	89.6
	CI-	2.4	3.4	1.1	3.6	1.0		CI-	89.2	91.2	86.5	90.6	87.2
Crack	%	0.3	0.5	0.1	0.5	0.2	Tobacco	%	60.0	61.6	58.4	57.5	62.3
	CI+	0.5	1.0	0.3	0.9	0.4		CI+	61.6	64.0	60.4	60.2	64.2
	CI-	0.2	0.2	0.0	0.2	0.1		CI-	58.4	59.1	56.4	54.9	60.3
Cocaine powder	%	2.9	4.1	1.6	4.5	1.4							
	CI+	3.5	5.2	2.2	5.8	1.9							
	CI-	2.3	3.2	1.1	3.5	0.9							
Amphetamines	%	3.0	4.0	1.9	4.8	1.3							
	CI+	3.6	5.1	2.6	6.1	1.8							
	CI-	2.4	3.1	1.3	3.7	0.9							
Ecstasy	%	3.7	4.9	2.6	7.1	0.7							
	CI+	4.4	6.1	3.3	8.6	1.1							
	CI-	3.1	3.8	1.9	5.8	0.4							

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

All HB Areas

Table 6.2: Ireland – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	5.6	7.8	3.4
	CI+	6.5	9.3	4.2
	CI-	4.8	6.5	2.6
Cannabis	%	5.0	7.2	2.9
	CI+	5.9	8.6	3.7
	CI-	4.3	5.9	2.2
Heroin	%	0.1	0.1	0.1
	CI+	0.3	0.3	0.4
	CI-	0.0	0.0	0.0
Methadone	%	0.2	0.2	0.2
	CI+	0.4	0.5	0.5
	CI-	0.1	0.0	0.1
Other opiates <sup>2</sup>	%	0.5	0.4	0.5
	CI+	0.8	0.9	0.9
	CI-	0.3	0.2	0.3
Cocaine (total including crack)	%	1.1	1.7	0.5
	CI+	1.5	2.5	0.8
	CI-	0.7	1.1	0.2
Crack	%	0.1	0.1	0.1
	CI+	0.2	0.3	0.3
	CI-	0.0	0.0	0.0
Cocaine powder	%	1.1	1.7	0.5
	CI+	1.5	2.5	0.8
	CI-	0.7	1.1	0.2
Amphetamines	%	0.4	0.7	0.2
	CI+	0.7	1.2	0.4
	CI-	0.2	0.3	0.1
Ecstasy	%	1.1	1.5	0.6
	CI+	1.5	2.3	1.0
	CI-	0.8	1.0	0.3

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.1	0.2	0.0
	CI+	0.2	0.5	0.1
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.4	0.6	0.1
	CI+	0.6	1.2	0.3
	CI-	0.2	0.3	0.0
Solvents	%	0.1	0.0	0.1
	CI+	0.2	0.3	0.3
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.4	0.4	0.8
	CI+	0.6	0.9	1.3
	CI-	0.2	0.2	0.4
Sedatives, tranquillisers, anti-depressants	%	5.7	4.2	3.7
	CI+	6.5	5.2	4.7
	CI-	5.0	3.3	2.9
Alcohol	%	83.8	86.0	86.5
	CI+	85.0	87.6	88.2
	CI-	82.7	84.4	84.7
Tobacco	%	38.0	38.2	43.2
	CI+	39.7	40.7	45.9
	CI-	36.4	35.8	40.5

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

All HB Areas

Table 6.3: Ireland – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	3.0	4.1	1.7
	CI+	3.6	5.3	2.4
	CI-	2.4	3.1	1.2
Cannabis	%	2.6	3.4	1.7
	CI+	3.2	4.5	2.3
	CI-	2.0	2.5	1.2
Heroin	%	0.1	0.1	0.1
	CI+	0.2	0.3	0.3
	CI-	0.0	0.0	0.0
Methadone	%	0.1	0.1	0.2
	CI+	0.3	0.3	0.5
	CI-	0.1	0.0	0.1
Other opiates <sup>2</sup>	%	0.2	0.2	0.1
	CI+	0.3	0.5	0.4
	CI-	0.1	0.0	0.0
Cocaine (total including crack)	%	0.3	0.7	0.0
	CI+	0.6	1.2	0.1
	CI-	0.2	0.3	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.1	0.2	0.1
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.3	0.7	0.0
	CI+	0.6	1.2	0.1
	CI-	0.2	0.3	0.0
Amphetamines	%	0.2	0.2	0.1
	CI+	0.3	0.6	0.3
	CI-	0.1	0.1	0.0
Ecstasy	%	0.3	0.6	0.0
	CI+	0.5	1.0	0.2
	CI-	0.2	0.3	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.1	0.3	0.1
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.1	0.1	0.1
	CI+	0.2	0.4	0.4
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.1
	CI+	0.1	0.3	0.3
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.1	0.1	0.2
	CI+	0.2	0.4	0.5
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	4.0	3.0	2.1
	CI+	4.6	4.0	2.9
	CI-	3.4	2.3	1.5
Alcohol	%	74.0	78.4	74.6
	CI+	75.4	80.3	76.8
	CI-	72.5	76.4	72.4
Tobacco	%	33.2	33.5	37.0
	CI+	34.9	35.9	39.8
	CI-	31.6	31.1	34.3

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 7 – Ireland

East Coast Area Health Board (ECAHB)

Table 7.1: ECAHB – Lifetime Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males			15-64	Males		
Any illegal drugs <sup>1</sup>	%	25.9	34.0	17.6	35.9	18.0	LSD	%
	CI+	31.7	44.0	24.3	45.0	24.4		CI+
	CI-	20.7	24.9	12.1	27.5	12.7		CI-
Cannabis	%	24.5	32.3	16.4	33.4	17.4	Magic mushrooms	%
	CI+	30.0	41.7	23.2	41.5	23.9		CI+
	CI-	19.5	23.9	10.8	25.9	12.1		CI-
Heroin	%	0.7	0.9	0.4	1.1	0.3	Solvents	%
	CI+	2.1	3.7	2.5	4.7	1.9		CI+
	CI-	0.1	0.1	0.0	0.1	0.0		CI-
Methadone	%	0.5	0.5	0.4	1.1	0.0	Poppers <sup>3</sup>	%
	CI+	1.9	3.5	2.5	4.7	1.3		CI+
	CI-	0.1	0.0	0.0	0.1	0.0		CI-
Other opiates <sup>2</sup>	%	3.8	2.6	5.1	3.2	4.3	Sedatives, tranquillisers, anti-depressants	%
	CI+	7.9	6.2	10.5	8.3	9.8		CI+
	CI-	1.5	0.8	2.0	0.8	1.4		CI-
Cocaine (total including crack)	%	6.3	10.5	2.0	10.5	3.0	Alcohol	%
	CI+	9.9	16.3	5.1	16.8	6.1		CI+
	CI-	3.7	6.3	0.5	6.0	1.2		CI-
Crack	%	1.0	1.5	0.4	1.8	0.3	Tobacco	%
	CI+	2.6	4.8	2.5	6.0	1.9		CI+
	CI-	0.3	0.3	0.0	0.3	0.0		CI-
Cocaine powder	%	6.3	10.5	2.0	10.5	3.0		
	CI+	9.9	16.3	5.1	16.8	6.1		
	CI-	3.7	6.3	0.5	6.0	1.2		
Amphetamines	%	4.8	6.6	3.0	8.0	2.3		
	CI+	7.6	11.1	5.9	14.4	4.9		
	CI-	2.9	3.6	1.2	3.8	0.9		
Ecstasy	%	5.4	7.8	3.0	9.8	1.9		
	CI+	8.3	12.3	5.9	16.0	4.5		
	CI-	3.3	4.5	1.2	5.3	0.6		

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## East Coast Area Health Board (ECAHB)

Table 7.2: ECAHB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	6.3	10.0	2.5
	CI+	9.4	15.5	5.3
	CI-	4.0	5.9	1.0
Cannabis	%	6.1	9.6	2.5
	CI+	9.0	14.9	5.3
	CI-	3.9	5.7	1.0
Heroin	%	0.2	0.0	0.4
	CI+	1.3	1.8	2.5
	CI-	0.0	0.0	0.0
Methadone	%	0.2	0.0	0.4
	CI+	1.3	1.8	2.5
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.4	0.0	0.8
	CI+	1.5	1.8	2.9
	CI-	0.0	0.0	0.1
Cocaine (total including crack)	%	2.3	4.2	0.4
	CI+	5.1	9.3	2.5
	CI-	0.8	1.4	0.0
Crack	%	0.5	0.6	0.4
	CI+	2.0	3.8	2.5
	CI-	0.1	0.0	0.0
Cocaine powder	%	2.3	4.2	0.4
	CI+	5.1	9.3	2.5
	CI-	0.8	1.4	0.0
Amphetamines	%	0.8	1.6	0.0
	CI+	2.4	5.0	1.5
	CI-	0.1	0.3	0.0
Ecstasy	%	2.5	4.0	0.9
	CI+	4.7	7.9	3.1
	CI-	1.1	1.7	0.1

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.8	1.8	2.2
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.8	1.8	2.2
	CI-	0.0	0.0	0.0
Solvents	%	0.2	0.0	0.4
	CI+	1.3	1.8	2.4
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	1.0	1.5	0.4
	CI+	2.6	4.8	2.4
	CI-	0.3	0.3	0.0
Sedatives, tranquillisers, anti-depressants	%	6.9	6.2	7.6
	CI+	10.4	10.8	12.6
	CI-	4.3	3.2	4.1
Alcohol	%	89.2	91.8	86.5
	CI+	92.2	95.5	90.6
	CI-	85.5	86.3	81.5
Tobacco	%	39.2	39.9	38.6
	CI+	43.9	48.3	45.0
	CI-	34.7	31.9	32.4

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**East Coast Area Health Board (ECAHB)**

Table 7.3: ECAHB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64		
	15-64	Males				
Any illegal drugs <sup>1</sup>	%	4.1	6.0	2.2	8.1	1.0
	CI+	6.6	10.6	4.9	14.1	2.9
	CI-	2.4	3.0	0.7	4.1	0.2
Cannabis	%	3.8	5.4	2.2	7.5	1.0
	CI+	6.3	10.2	4.9	13.5	2.9
	CI-	2.1	2.5	0.7	3.5	0.2
Heroin	%	0.0	0.0	0.0	0.0	0.0
	CI+	0.8	1.8	1.5	2.2	1.3
	CI-	0.0	0.0	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0	0.0	0.0
	CI+	0.8	1.8	1.5	2.2	1.3
	CI-	0.0	0.0	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.0	0.0	0.0	0.0	0.0
	CI+	0.8	1.8	1.5	2.2	1.3
	CI-	0.0	0.0	0.0	0.0	0.0
Cocaine (total including crack)	%	0.5	0.9	0.0	0.6	0.3
	CI+	1.8	3.8	1.5	4.2	1.9
	CI-	0.0	0.1	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0	0.0	0.0
	CI+	0.8	1.8	1.5	2.2	1.3
	CI-	0.0	0.0	0.0	0.0	0.0
Cocaine powder	%	0.5	0.9	0.0	0.6	0.3
	CI+	1.8	3.8	1.5	4.2	1.9
	CI-	0.0	0.1	0.0	0.0	0.0
Amphetamines	%	0.5	1.0	0.0	0.7	0.3
	CI+	1.9	4.0	1.5	4.7	1.9
	CI-	0.0	0.1	0.0	0.0	0.0
Ecstasy	%	0.9	1.7	0.0	1.9	0.0
	CI+	2.5	5.2	1.5	6.3	1.3
	CI-	0.2	0.3	0.0	0.3	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.5
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.5
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.5
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.5
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	4.9	4.5	5.3
	CI+	7.6	8.3	10.2
	CI-	3.0	2.1	2.3
Alcohol	%	81.2	86.6	75.6
	CI+	85.6	92.1	81.4
	CI-	76.1	79.3	69.0
Tobacco	%	32.0	33.4	30.5
	CI+	37.6	43.1	36.6
	CI-	26.7	24.6	24.9

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 8 – Ireland

Midland Health Board (MHB)

Table 8.1: MHB – Lifetime Prevalence (%)

	All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		
Any illegal drugs <sup>1</sup>	%	11.0	13.3	8.5	15.2	7.4	LSD	%	1.6	2.6	0.6	2.6	0.8
	CI+	15.9	20.1	15.5	24.6	12.4		CI+	3.7	6.8	3.2	6.7	4.1
	CI-	7.2	8.1	4.0	8.3	4.0		CI-	0.5	0.6	0.0	0.7	0.0
Cannabis	%	10.7	13.3	7.9	14.6	7.4	Magic mushrooms	%	1.8	2.8	0.7	2.9	0.8
	CI+	15.7	20.1	14.9	24.1	12.4		CI+	4.1	7.1	3.3	7.9	4.1
	CI-	6.8	8.1	3.5	7.8	4.0		CI-	0.6	0.7	0.0	0.6	0.0
Heroin	%	0.3	0.6	0.0	0.6	0.0	Solvents	%	1.2	1.7	0.6	2.6	0.0
	CI+	1.7	3.7	2.1	3.8	2.1		CI+	3.1	5.6	3.2	6.7	2.1
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	0.3	0.3	0.0	0.7	0.0
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%	1.6	2.5	0.7	3.5	0.0
	CI+	1.2	2.6	2.1	2.6	2.1		CI+	3.7	7.2	3.3	8.0	2.1
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	0.5	0.5	0.0	1.1	0.0
Other opiates <sup>2</sup>	%	1.3	1.5	1.1	1.3	1.3	Sedatives, tranquillisers, anti-depressants	%	10.5	9.5	11.4	3.9	16.0
	CI+	3.3	5.4	4.0	4.8	4.3		CI+	14.4	15.7	17.0	8.5	22.3
	CI-	0.4	0.2	0.1	0.1	0.2		CI-	7.3	5.2	7.1	1.3	10.9
Cocaine (total including crack)	%	1.3	2.0	0.6	1.9	0.8	Alcohol	%	87.1	92.6	81.3	89.5	85.1
	CI+	3.3	6.0	3.2	5.8	4.1		CI+	90.6	96.7	86.7	94.1	90.0
	CI-	0.4	0.4	0.0	0.4	0.0		CI-	82.9	86.2	74.8	83.3	78.9
Crack	%	0.3	0.6	0.0	0.6	0.0	Tobacco	%	60.4	61.5	59.2	55.5	64.6
	CI+	1.7	3.7	2.1	3.8	2.1		CI+	66.3	69.7	66.5	64.2	71.7
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	54.3	52.9	51.6	46.5	57.0
Cocaine powder	%	1.3	2.0	0.6	1.9	0.8							
	CI+	3.3	6.0	3.2	5.8	4.1							
	CI-	0.4	0.4	0.0	0.4	0.0							
Amphetamines	%	0.6	1.2	0.0	1.3	0.0							
	CI+	2.2	4.7	2.1	4.8	2.1							
	CI-	0.1	0.1	0.0	0.1	0.0							
Ecstasy	%	2.0	2.6	1.3	3.4	0.8							
	CI+	4.2	6.8	4.3	8.0	4.1							
	CI-	0.7	0.6	0.2	1.0	0.0							

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Midland Health Board (MHB)**

Table 8.2: MHB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	2.8	3.6	1.9
	CI+	5.5	8.2	5.8
	CI-	1.2	1.2	0.4
Cannabis	%	2.8	3.6	1.9
	CI+	5.5	8.2	5.8
	CI-	1.2	1.2	0.4
Heroin	%	0.0	0.0	0.0
	CI+	1.2	2.6	2.1
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	1.2	2.6	2.1
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.0	0.0	0.0
	CI+	1.2	2.6	2.1
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.3	0.6	0.0
	CI+	1.7	3.7	2.1
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	1.2	2.6	2.1
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.3	0.6	0.0
	CI+	1.7	3.7	2.1
	CI-	0.0	0.0	0.0
Amphetamines	%	0.3	0.6	0.0
	CI+	1.7	3.7	2.1
	CI-	0.0	0.0	0.0
Ecstasy	%	0.9	1.2	0.7
	CI+	2.7	4.7	3.3
	CI-	0.2	0.1	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	1.2	2.6	2.1
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.3	0.6	0.0
	CI+	1.7	3.7	2.1
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	1.2	2.6	2.1
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.0	0.0	0.0
	CI+	1.2	2.6	2.1
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	4.2	3.6	0.6
	CI+	7.1	9.4	3.8
	CI-	2.3	0.9	0.0
Alcohol	%	78.6	83.5	82.1
	CI+	83.0	90.0	88.0
	CI-	73.7	74.9	74.7
Tobacco	%	36.5	35.5	29.7
	CI+	42.4	44.1	37.1
	CI-	30.9	27.5	35.4

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Midland Health Board (MHB)**

Table 8.3: MHB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	1.0	2.0	0.0
CI+	2.9	6.0	2.1	4.8
CI-	0.2	0.4	0.0	0.1
Cannabis	%	1.0	2.0	0.0
CI+	2.9	6.0	2.1	4.8
CI-	0.2	0.4	0.0	0.1
Heroin	%	0.0	0.0	0.0
CI+	1.2	2.6	2.1	2.6
CI-	0.0	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
CI+	1.2	2.6	2.1	2.6
CI-	0.0	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.0	0.0	0.0
CI+	1.2	2.6	2.1	2.6
CI-	0.0	0.0	0.0	0.0
Cocaine (total including crack)	%	0.3	0.6	0.0
CI+	1.7	3.7	2.1	3.8
CI-	0.0	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
CI+	1.2	2.6	2.1	2.6
CI-	0.0	0.0	0.0	0.0
Cocaine powder	%	0.3	0.6	0.0
CI+	1.7	3.7	2.1	3.8
CI-	0.0	0.0	0.0	0.0
Amphetamines	%	0.0	0.0	0.0
CI+	1.2	2.6	2.1	2.6
CI-	0.0	0.0	0.0	0.0
Ecstasy	%	0.0	0.0	0.0
CI+	1.2	2.6	2.1	2.6
CI-	0.0	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
CI+	1.2	2.6	2.1	2.6
CI-	0.0	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
CI+	1.2	2.6	2.1	2.6
CI-	0.0	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
CI+	1.2	2.6	2.1	2.6
CI-	0.0	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.0	0.0	0.0
CI+	1.2	2.6	2.1	2.6
CI-	0.0	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	2.8	2.9	2.6
CI+	5.3	8.8	6.2	2.6
CI-	1.3	0.5	0.8	0.0
Alcohol	%	69.0	73.9	71.0
CI+	75.8	82.2	70.9	79.3
CI-	61.5	64.2	56.3	61.6
Tobacco	%	31.4	30.8	38.7
CI+	37.0	39.2	47.9	32.4
CI-	26.2	23.2	25.3	30.1

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 9 – Ireland

Mid-Western Health Board (MWHB)

Table 9.1: MWHB – Lifetime Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males			15-64	Males		
Any illegal drugs <sup>1</sup>	%	12.0	15.3	8.5	17.8	7.0	LSD	%
	CI+	18.0	26.2	13.4	28.6	10.7		CI+
	CI-	7.4	7.7	5.0	9.8	4.2		CI-
Cannabis	%	10.9	13.7	8.1	16.5	6.1	Magic mushrooms	%
	CI+	15.9	23.2	12.6	26.5	9.6		CI+
	CI-	7.1	7.0	4.7	9.2	3.5		CI-
Heroin	%	0.0	0.0	0.0	0.0	0.0	Solvents	%
	CI+	0.9	2.0	1.5	2.3	1.4		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%
	CI+	0.9	2.0	1.5	2.3	1.4		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Other opiates <sup>2</sup>	%	1.8	1.2	2.5	2.2	1.5	Sedatives, tranquillisers, anti-depressants	%
	CI+	3.6	4.5	5.4	6.1	3.8		CI+
	CI-	0.8	0.1	0.9	0.5	0.4		CI-
Cocaine (total including crack)	%	1.1	1.2	1.1	1.9	0.5	Alcohol	%
	CI+	3.4	7.1	3.5	5.6	2.3		CI+
	CI-	0.2	0.0	0.2	0.4	0.0		CI-
Crack	%	0.0	0.0	0.0	0.0	0.0	Tobacco	%
	CI+	0.9	2.0	1.5	2.3	1.4		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Cocaine powder	%	1.1	1.2	1.1	1.9	0.5		
	CI+	3.4	7.1	3.5	5.6	2.3		
	CI-	0.2	0.0	0.2	0.4	0.0		
Amphetamines	%	1.3	1.2	1.5	2.3	0.5		
	CI+	3.0	4.4	4.0	6.1	2.4		
	CI-	0.5	0.1	0.4	0.6	0.0		
Ecstasy	%	1.7	0.7	2.7	3.7	0.0		
	CI+	3.9	4.2	6.6	8.9	1.4		
	CI-	0.6	0.0	0.8	1.1	0.0		

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Mid-Western Health Board (MWHB)**

Table 9.2: MWHB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	3.2	3.9	2.5
	CI+	8.2	11.3	6.2
	CI-	0.8	0.8	0.7
Cannabis	%	3.0	3.9	2.0
	CI+	7.3	11.3	4.8
	CI-	0.8	0.8	0.6
Heroin	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.6	0.7	0.6
	CI+	2.3	4.2	3.1
	CI-	0.1	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.6	0.7	0.6
	CI+	2.3	4.2	3.1
	CI-	0.1	0.0	0.0
Amphetamines	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Ecstasy	%	0.6	0.7	0.5
	CI+	2.1	4.2	2.4
	CI-	0.1	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	4.0	4.4	3.6
	CI+	7.1	9.8	7.1
	CI-	2.0	1.5	1.5
Alcohol	%	81.2	82.6	79.7
	CI+	85.5	88.6	84.6
	CI-	76.1	75.1	74.0
Tobacco	%	35.6	35.0	36.3
	CI+	41.6	43.5	44.1
	CI-	30.0	27.1	28.9

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Mid-Western Health Board (MWHB)**

Table 9.3: MWHB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs	%	1.6	1.9	1.4
	CI+	4.9	7.4	3.8
	CI-	0.3	0.2	0.3
Cannabis	%	1.6	1.9	1.4
	CI+	4.9	7.4	3.8
	CI-	0.3	0.2	0.3
Heroin	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Other opiates	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Amphetamines	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Ecstasy	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Poppers	%	0.0	0.0	0.0
	CI+	0.9	2.0	1.5
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	3.3	3.3	0.6
	CI+	6.6	8.8	4.1
	CI-	1.4	0.8	0.0
Alcohol	%	71.8	74.0	72.7
	CI+	77.5	81.5	80.2
	CI-	65.6	65.4	64.1
Tobacco	%	29.7	30.3	32.6
	CI+	35.3	39.4	40.5
	CI-	24.6	22.2	25.3

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 10 – Ireland

North Eastern Health Board (NEHB)

Table 10.1: NEHB – Lifetime Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males			15-64	Males		
Any illegal drugs <sup>1</sup>	%	18.9	24.7	12.9	32.4	7.2	LSD	%
	CI+	24.3	33.7	17.7	41.9	12.9		CI+
	CI-	14.3	17.0	9.1	23.9	3.5		CI-
Cannabis	%	17.8	24.2	11.0	30.4	6.7	Magic mushrooms	%
	CI+	23.1	33.2	15.6	40.2	12.4		CI+
	CI-	13.2	16.5	7.4	21.8	3.1		CI-
Heroin	%	0.4	0.8	0.0	0.9	0.0	Solvents	%
	CI+	2.2	4.9	1.5	5.2	1.4		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%
	CI+	0.8	2.0	1.5	2.1	1.4		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Other opiates <sup>2</sup>	%	1.3	0.5	2.1	0.8	1.7	Sedatives, tranquillisers, anti-depressants	%
	CI+	2.9	3.1	5.1	3.5	4.1		CI+
	CI-	0.5	0.0	0.7	0.1	0.5		CI-
Cocaine (total including crack)	%	1.2	2.4	0.0	2.1	0.5	Alcohol	%
	CI+	3.0	6.4	1.5	6.2	2.4		CI+
	CI-	0.4	0.6	0.0	0.4	0.0		CI-
Crack	%	0.3	0.6	0.0	0.6	0.0	Tobacco	%
	CI+	1.6	3.5	1.5	3.6	1.4		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Cocaine powder	%	0.9	1.8	0.0	1.5	0.5		
	CI+	2.7	5.8	1.5	5.6	2.4		
	CI-	0.2	0.3	0.0	0.1	0.0		
Amphetamines	%	2.6	3.8	1.4	4.0	1.4		
	CI+	4.6	8.1	3.8	8.4	3.7		
	CI-	1.3	1.4	0.3	1.5	0.3		
Ecstasy	%	2.6	3.2	1.9	4.9	0.5		
	CI+	5.2	8.6	4.5	10.9	2.4		
	CI-	1.0	0.7	0.6	1.7	0.0		

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## North Eastern Health Board (NEHB)

Table 10.2: NEHB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	6.4	8.3	4.3
	CI+	9.9	13.9	8.1
	CI-	3.8	4.5	2.0
Cannabis	%	5.2	7.8	2.4
	CI+	8.2	12.9	5.7
	CI-	3.0	4.2	0.7
Heroin	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.2	0.0	0.4
	CI+	1.2	2.0	2.2
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Amphetamines	%	0.2	0.0	0.5
	CI+	1.4	2.0	2.5
	CI-	0.0	0.0	0.0
Ecstasy	%	0.5	0.0	1.0
	CI+	1.7	2.0	3.2
	CI-	0.1	0.0	0.2
				1.1
				0.0

	All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64
LSD	%	0.0	0.0	0.0	0.0
	CI+	0.8	2.0	1.5	2.1
	CI-	0.0	0.0	0.0	0.0
Magic mushrooms	%	0.5	1.0	0.0	0.6
	CI+	1.8	3.8	1.5	3.7
	CI-	0.1	0.1	0.0	0.0
Solvents	%	0.5	0.6	0.5	1.1
	CI+	1.9	3.5	2.4	4.2
	CI-	0.1	0.0	0.0	0.1
Poppers <sup>3</sup>	%	1.2	0.6	1.9	2.6
	CI+	2.8	3.5	4.5	6.2
	CI-	0.4	0.0	0.6	0.8
Sedatives, tranquillisers, anti-depressants	%	2.1	1.4	2.8	1.1
	CI+	3.9	5.3	5.7	4.0
	CI-	0.9	0.1	1.1	0.1
Alcohol	%	82.3	84.5	80.0	85.1
	CI+	85.8	89.9	84.8	90.7
	CI-	78.4	77.6	74.5	77.8
Tobacco	%	34.6	37.1	31.9	43.0
	CI+	40.9	46.8	39.2	52.3
	CI-	28.6	28.2	25.1	34.0
				27.3	34.1
				21.1	21.1

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**North Eastern Health Board (NEHB)**

Table 10.3: NEHB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	2.4	3.4	1.4
	CI+	5.4	8.8	3.8
	CI-	0.8	0.9	0.3
Cannabis	%	1.9	2.9	0.9
	CI+	4.3	8.4	3.0
	CI-	0.7	0.6	0.1
Heroin	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Amphetamines	%	0.2	0.0	0.5
	CI+	1.4	2.0	2.5
	CI-	0.0	0.0	0.0
Ecstasy	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.8	2.0	1.5
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.2	0.4	0.0
	CI+	1.3	2.8	1.5
	CI-	0.0	0.0	0.0
Solvents	%	0.3	0.6	0.6
	CI+	1.6	3.5	1.5
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.2	0.0	0.5
	CI+	1.4	2.0	3.1
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	1.6	0.9	0.5
	CI+	3.3	3.7	3.3
	CI-	0.6	0.1	0.0
Alcohol	%	71.2	77.7	72.2
	CI+	75.7	84.3	79.2
	CI-	66.4	69.9	64.3
Tobacco	%	31.1	33.7	36.9
	CI+	37.4	42.9	46.4
	CI-	25.2	25.3	28.0

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 11 – Ireland

North Western Health Board (NWHB)

Table 11.1: NWHB – Lifetime Prevalence (%)

	All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		
Any illegal drugs <sup>1</sup>	%	10.6	12.7	8.5	13.3	8.6	LSD	%	0.3	0.6	0.0	0.0	0.5
	CI+	14.8	19.4	13.6	20.5	13.3		CI+	1.7	3.5	2.0	2.9	2.7
	CI-	7.3	7.7	4.9	7.9	5.1		CI-	0.0	0.0	0.0	0.0	0.0
Cannabis	%	9.3	11.9	6.6	10.7	8.2	Magic mushrooms	%	1.5	2.0	1.1	2.2	1.0
	CI+	13.2	18.2	11.7	17.4	12.9		CI+	3.6	6.9	3.9	7.7	3.5
	CI-	6.2	7.2	3.2	5.9	4.8		CI-	0.5	0.2	0.1	0.3	0.1
Heroin	%	0.0	0.0	0.0	0.0	0.0	Solvents	%	0.7	0.0	1.4	1.0	0.4
	CI+	1.1	2.4	2.0	2.9	1.8		CI+	2.3	2.4	4.3	5.4	2.5
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	0.1	0.0	0.2	0.0	0.0
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%	0.9	1.2	0.6	1.3	0.5
	CI+	1.1	2.4	2.0	2.9	1.8		CI+	2.6	4.4	3.1	5.2	2.7
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	0.2	0.1	0.0	0.1	0.0
Other opiates <sup>2</sup>	%	1.2	0.0	2.4	1.7	0.8	Sedatives, tranquillisers, anti-depressants	%	10.6	9.0	12.3	7.4	13.2
	CI+	3.5	2.4	6.6	6.0	3.2		CI+	15.2	14.7	18.5	13.7	18.6
	CI-	0.2	0.0	0.6	0.2	0.1		CI-	7.0	4.9	7.6	3.4	8.9
Cocaine (total including crack)	%	0.0	0.0	0.0	0.0	0.0	Alcohol	%	86.0	90.8	81.0	88.3	84.2
	CI+	1.1	2.4	2.0	2.9	1.8		CI+	90.8	94.9	88.4	94.2	89.1
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	79.9	85.1	71.7	79.5	78.2
Crack	%	0.0	0.0	0.0	0.0	0.0	Tobacco	%	58.2	61.1	55.3	48.3	66.0
	CI+	1.1	2.4	2.0	2.9	1.8		CI+	64.4	69.6	64.4	58.9	72.5
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	51.9	52.1	45.8	37.9	59.1
Cocaine powder	%	0.0	0.0	0.0	0.0	0.0							
	CI+	1.1	2.4	2.0	2.9	1.8							
	CI-	0.0	0.0	0.0	0.0	0.0							
Amphetamines	%	0.3	0.6	0.0	0.0	0.5							
	CI+	1.7	3.5	2.0	2.9	2.7							
	CI-	0.0	0.0	0.0	0.0	0.0							
Ecstasy	%	0.3	0.0	0.6	0.6	0.0							
	CI+	1.6	2.4	3.1	4.1	1.8							
	CI-	0.0	0.0	0.0	0.0	0.0							

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**North Western Health Board (NWHB)**

Table 11.2: NWHB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	2.6	4.2	0.9
	CI+	5.6	9.9	3.6
	CI-	0.9	1.3	0.1
Cannabis	%	2.2	3.4	0.9
	CI+	4.4	7.7	3.6
	CI-	0.9	1.1	0.1
Heroin	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Amphetamines	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Ecstasy	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.4	0.8	0.0
	CI+	1.9	4.1	2.0
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	5.9	4.3	7.4
	CI+	9.2	8.9	12.3
	CI-	3.5	1.7	4.1
Alcohol	%	77.0	82.0	71.9
	CI+	83.4	87.8	81.5
	CI-	69.5	74.9	60.5
Tobacco	%	32.8	33.6	32.0
	CI+	39.0	41.7	39.9
	CI-	27.0	26.1	24.8

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**North Western Health Board (NWHB)**

Table 11.3: NWHB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	0.2	0.0	0.5
	CI+	1.6	2.4	2.9
	CI-	0.0	0.0	0.0
Cannabis	%	0.2	0.0	0.5
	CI+	1.6	2.4	2.9
	CI-	0.0	0.0	0.0
Heroin	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Amphetamines	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0
Ecstasy	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.0
	CI-	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.9
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.9
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.9
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.0	0.0	0.0
	CI+	1.1	2.4	2.9
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	4.5	3.6	5.4
	CI+	7.3	8.0	9.8
	CI-	2.5	1.3	2.6
Alcohol	%	67.6	74.8	60.3
	CI+	73.0	81.5	69.8
	CI-	61.9	67.1	50.1
Tobacco	%	28.5	28.9	28.1
	CI+	33.7	36.8	35.3
	CI-	23.7	21.8	21.7

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 12 – Ireland

Northern Area Health Board (NAHB)

Table 12.1: NAHB – Lifetime Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	29.5	40.3	19.6
	CI+	36.0	50.3	26.4
	CI-	23.6	30.8	13.8
Cannabis	%	26.9	35.2	19.1
	CI+	32.8	44.3	26.2
	CI-	21.5	26.8	13.3
Heroin	%	0.9	1.2	0.7
	CI+	2.3	3.8	2.6
	CI-	0.3	0.2	0.1
Methadone	%	1.0	1.2	0.8
	CI+	2.8	3.9	4.5
	CI-	0.2	0.2	0.0
Other opiates <sup>2</sup>	%	8.4	6.6	10.0
	CI+	11.7	11.2	15.1
	CI-	5.8	3.5	6.2
Cocaine (total including crack)	%	5.2	8.1	2.5
	CI+	8.1	13.5	6.9
	CI-	3.1	4.3	0.6
Crack	%	0.6	1.2	0.0
	CI+	1.9	4.1	1.2
	CI-	0.1	0.2	0.0
Cocaine powder	%	5.0	7.7	2.5
	CI+	8.0	13.2	6.9
	CI-	2.9	4.0	0.6
Amphetamines	%	3.8	6.0	1.7
	CI+	7.0	12.0	6.8
	CI-	1.8	2.5	0.1
Ecstasy	%	6.5	10.0	3.2
	CI+	10.0	17.0	7.4
	CI-	3.9	5.2	1.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	4.2	6.0	2.5
	CI+	7.4	11.7	7.1
	CI-	2.1	2.6	0.6
Magic mushrooms	%	5.1	8.2	2.3
	CI+	7.8	13.6	5.1
	CI-	3.2	4.4	0.8
Solvents	%	3.4	4.5	2.5
	CI+	5.8	9.4	5.6
	CI-	1.8	1.7	0.8
Poppers <sup>3</sup>	%	4.0	6.1	2.0
	CI+	7.6	12.6	5.1
	CI-	1.8	2.3	0.5
Sedatives, tranquillisers, anti-depressants	%	17.5	12.9	21.7
	CI+	21.0	18.9	26.9
	CI-	14.3	8.2	17.2
Alcohol	%	93.5	94.8	92.3
	CI+	95.7	97.5	95.3
	CI-	90.7	90.6	88.3
Tobacco	%	62.5	61.8	63.1
	CI+	67.7	70.5	69.0
	CI-	57.0	52.5	56.9

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Northern Area Health Board (NAHB)**

Table 12.2: NAHB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	8.4	13.3	3.9
	CI+	12.0	20.3	8.2
	CI-	5.6	8.0	1.5
Cannabis	%	7.7	12.2	3.4
	CI+	11.1	19.0	7.9
	CI-	5.0	7.2	1.1
Heroin	%	0.2	0.0	0.5
	CI+	1.4	1.6	2.5
	CI-	0.0	0.0	0.0
Methadone	%	0.4	0.3	0.5
	CI+	1.6	2.3	2.5
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.7	1.1	0.3
	CI+	2.1	4.4	1.7
	CI-	0.1	0.1	0.0
Cocaine (total including crack)	%	1.7	2.7	0.8
	CI+	4.0	7.0	2.9
	CI-	0.6	0.7	0.1
Crack	%	0.0	0.0	0.0
	CI+	0.7	1.6	1.2
	CI-	0.0	0.0	0.0
Cocaine powder	%	1.7	2.7	0.8
	CI+	4.0	7.0	2.9
	CI-	0.6	0.7	0.1
Amphetamines	%	0.3	0.5	0.0
	CI+	1.6	3.5	1.2
	CI-	0.0	0.0	0.0
Ecstasy	%	1.6	2.7	0.5
	CI+	3.4	6.6	2.5
	CI-	0.6	0.8	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.7	1.6	1.2
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.9	1.6	0.3
	CI+	2.5	5.2	1.7
	CI-	0.2	0.3	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.7	1.6	1.2
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.3	0.5	0.0
	CI+	1.6	3.5	1.2
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	8.9	5.2	12.3
	CI+	11.7	9.8	16.6
	CI-	6.6	2.3	8.8
Alcohol	%	88.8	92.0	85.9
	CI+	91.8	95.2	89.8
	CI-	85.2	87.6	81.1
Tobacco	%	42.9	40.8	44.9
	CI+	49.4	50.7	52.2
	CI-	36.6	31.5	37.7

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Northern Area Health Board (NAHB)**

Table 12.3: NAHB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	5.3	8.5	2.3
	CI+	8.5	14.7	6.9
	CI-	3.1	4.4	0.5
Cannabis	%	4.5	6.9	2.3
	CI+	7.5	12.3	6.9
	CI-	2.5	3.4	0.5
Heroin	%	0.2	0.0	0.5
	CI+	1.4	1.6	2.5
	CI-	0.0	0.0	0.0
Methadone	%	0.4	0.3	0.5
	CI+	1.6	2.3	2.5
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.3	0.5	0.0
	CI+	1.6	3.5	1.2
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.8	1.6	0.0
	CI+	2.4	5.3	1.2
	CI-	0.2	0.3	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.7	1.6	1.2
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.8	1.6	0.0
	CI+	2.4	5.3	1.2
	CI-	0.2	0.3	0.0
Amphetamines	%	0.3	0.5	0.0
	CI+	1.6	3.5	1.2
	CI-	0.0	0.0	0.0
Ecstasy	%	0.3	0.5	0.0
	CI+	1.6	3.5	1.2
	CI-	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.7	1.6	1.2
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.7	1.6	1.2
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.7	1.6	1.2
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.3	0.5	0.5
	CI+	1.6	3.5	1.2
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	5.6	3.1	2.7
	CI+	7.9	7.2	6.0
	CI-	3.8	1.0	0.9
Alcohol	%	80.7	86.9	82.7
	CI+	84.5	91.0	87.5
	CI-	76.5	81.8	77.0
Tobacco	%	38.3	37.7	41.5
	CI+	44.6	47.6	52.0
	CI-	32.3	28.5	31.5

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 13 – Ireland

South Eastern Health Board (SEHB)

Table 13.1: SEHB – Lifetime Prevalence (%)

	All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		
Any illegal drugs <sup>1</sup>	%	18.5	24.8	12.0	27.2	11.4	LSD	%	4.6	7.2	1.9	8.5	1.4
	CI+	22.9	31.9	15.9	35.2	15.2		CI+	6.7	11.7	3.9	13.5	3.1
	CI-	14.6	18.5	8.8	20.1	8.2		CI-	3.0	4.0	0.7	4.9	0.5
Cannabis	%	16.8	22.0	11.4	25.4	9.8	Magic mushrooms	%	5.7	9.5	1.8	9.3	2.7
	CI+	21.0	28.9	15.2	33.4	13.1		CI+	8.1	14.1	4.7	14.9	4.9
	CI-	13.1	16.0	8.3	18.5	7.0		CI-	3.8	5.9	0.5	5.3	1.4
Heroin	%	0.0	0.0	0.0	0.0	0.0	Solvents	%	1.1	1.9	0.3	2.0	0.4
	CI+	0.6	1.4	1.1	1.7	0.9		CI+	2.4	4.8	1.6	5.4	1.7
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	0.4	0.5	0.0	0.5	0.0
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%	3.3	4.8	1.7	6.6	0.5
	CI+	0.6	1.4	1.1	1.7	0.9		CI+	5.4	9.3	3.9	11.7	1.8
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	1.8	2.0	0.6	3.3	0.1
Other opiates <sup>2</sup>	%	1.0	1.4	0.6	1.2	0.9	Sedatives, tranquillisers, anti-depressants	%	14.0	8.6	19.7	10.9	16.6
	CI+	2.2	3.7	2.1	3.7	2.4		CI+	17.2	12.7	24.3	15.9	21.0
	CI-	0.4	0.4	0.1	0.2	0.2		CI-	11.3	5.5	15.6	7.1	12.8
Cocaine (total including crack)	%	2.5	3.4	1.5	4.2	1.1	Alcohol	%	89.5	89.8	89.1	90.0	89.0
	CI+	4.2	6.5	3.4	8.7	2.7		CI+	92.3	93.2	92.8	94.6	92.1
	CI-	1.3	1.6	0.5	1.6	0.3		CI-	86.0	85.5	84.3	83.6	85.2
Crack	%	0.3	0.7	0.0	0.0	0.6	Tobacco	%	60.5	65.7	55.2	64.5	57.4
	CI+	1.2	2.6	1.1	1.7	2.0		CI+	64.5	71.5	60.5	73.0	62.3
	CI-	0.0	0.1	0.0	0.0	0.1		CI-	56.5	59.6	49.8	55.2	52.3
Cocaine powder	%	2.3	3.1	1.5	4.2	0.8							
	CI+	4.0	6.1	3.4	8.7	2.3							
	CI-	1.2	1.3	0.5	1.6	0.2							
Amphetamines	%	4.4	5.6	3.2	7.4	2.0							
	CI+	6.8	10.0	5.6	12.5	4.1							
	CI-	2.7	2.8	1.6	3.9	0.8							
Ecstasy	%	4.3	6.5	2.1	9.3	0.2							
	CI+	6.5	11.1	4.2	14.6	1.3							
	CI-	2.7	3.3	0.8	5.5	0.0							

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## South Eastern Health Board (SEHB)

Table 13.2: SEHB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	o/o	6.9	9.4	4.3
	CI+	9.9	14.3	7.7
	CI-	4.6	5.7	2.2
Cannabis	%	5.8	7.9	3.6
	CI+	8.5	12.9	6.8
	CI-	3.7	4.4	1.6
Heroin	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.4	0.3	0.6
	CI+	1.3	2.0	2.1
	CI-	0.1	0.0	0.1
Cocaine (total including crack)	%	1.7	2.5	1.0
	CI+	3.5	5.5	3.0
	CI-	0.7	0.8	0.2
Crack	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Cocaine powder	%	1.7	2.5	1.0
	CI+	3.5	5.5	3.0
	CI-	0.7	0.8	0.2
Amphetamines	%	0.9	1.0	0.8
	CI+	2.4	3.6	2.4
	CI-	0.2	0.1	0.1
Ecstasy	%	1.3	2.5	0.0
	CI+	2.7	5.7	1.1
	CI-	0.5	0.8	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.3	0.5	0.6
	CI+	1.3	2.9	3.3
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.8	1.5	1.7
	CI+	2.6	5.6	6.5
	CI-	0.1	0.1	0.2
Solvents	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.7
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.7	1.0	1.6
	CI+	2.0	3.6	4.8
	CI-	0.2	0.1	0.3
Sedatives, tranquillisers, anti-depressants	%	7.6	4.5	5.4
	CI+	10.3	8.0	9.7
	CI-	5.3	2.3	2.6
Alcohol	%	83.7	84.7	85.6
	CI+	87.3	89.0	91.6
	CI-	79.5	79.6	77.6
Tobacco	%	38.1	38.8	49.0
	CI+	42.1	45.1	57.7
	CI-	34.2	32.8	40.3

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## South Eastern Health Board (SEHB)

Table 13.3: SEHB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	3.1	4.5	1.7
	CI+	5.1	8.7	3.9
	CI-	1.7	1.9	0.6
Cannabis	%	2.1	2.5	1.7
	CI+	3.6	5.7	3.9
	CI-	1.1	0.8	0.6
Heroin	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.1	0.0	0.3
	CI+	0.9	1.4	1.6
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Amphetamines	%	0.2	0.0	0.5
	CI+	1.2	1.4	2.2
	CI-	0.0	0.0	0.0
Ecstasy	%	1.3	2.5	0.0
	CI+	2.7	5.7	1.1
	CI-	0.5	0.8	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.3	0.5	0.6
	CI+	1.3	2.9	3.3
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	4.9	3.6	2.6
	CI+	7.5	7.0	9.3
	CI-	3.0	1.5	3.9
Alcohol	%	74.1	77.4	73.1
	CI+	78.5	82.4	80.9
	CI-	69.3	71.8	64.1
Tobacco	%	32.4	33.4	39.8
	CI+	36.3	39.5	48.6
	CI-	28.7	27.7	31.5

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 14 – Ireland

South Western Area Health Board (SWAHB)

Table 14.1: SWAHB – Lifetime Prevalence (%)

	All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		
Any illegal drugs <sup>1</sup>	%	24.0	31.5	16.8	29.8	16.5	LSD	%	4.9	7.6	2.3	6.9	2.3
	CI+	29.5	40.3	22.3	37.3	21.9		CI+	7.2	11.5	5.1	10.9	4.8
	CI-	19.1	23.6	12.2	23.0	12.0		CI-	3.2	4.7	0.8	4.0	0.9
Cannabis	%	23.2	30.2	16.5	28.3	16.5	Magic mushrooms	%	5.6	8.8	2.5	8.0	2.5
	CI+	28.4	38.4	21.9	35.4	21.9		CI+	8.5	13.5	5.7	13.1	4.9
	CI-	18.5	22.8	11.9	21.9	12.0		CI-	3.4	5.3	0.8	4.4	1.1
Heroin	%	1.8	2.4	1.2	2.6	0.8	Solvents	%	1.6	1.8	1.4	2.8	0.0
	CI+	3.4	6.1	3.1	5.1	2.8		CI+	3.3	4.8	3.4	6.0	1.2
	CI-	0.8	0.7	0.4	1.1	0.1		CI-	0.6	0.4	0.4	1.0	0.0
Methadone	%	1.2	1.1	1.2	1.9	0.3	Poppers <sup>3</sup>	%	3.0	4.5	1.5	4.5	1.0
	CI+	2.4	3.8	3.1	4.2	1.8		CI+	5.3	8.4	3.5	8.7	3.0
	CI-	0.5	0.2	0.4	0.6	0.0		CI-	1.5	2.1	0.5	2.0	0.2
Other opiates <sup>2</sup>	%	6.0	3.4	8.4	5.4	6.8	Sedatives, tranquillisers, anti-depressants	%	13.4	10.7	15.9	11.2	16.2
	CI+	9.4	7.7	13.7	11.1	10.8		CI+	16.7	15.1	20.9	15.9	21.2
	CI-	3.5	1.2	4.7	2.0	4.0		CI-	10.6	7.3	11.7	7.6	12.0
Cocaine (total including crack)	%	5.0	6.7	3.3	7.3	1.9	Alcohol	%	91.2	93.9	88.6	92.5	89.5
	CI+	7.8	11.8	6.1	12.2	5.0		CI+	93.5	97.0	91.8	95.7	92.7
	CI-	2.9	3.3	1.5	3.9	0.5		CI-	88.4	89.2	84.7	88.0	85.5
Crack	%	0.4	0.5	0.3	0.7	0.0	Tobacco	%	63.8	66.7	60.9	62.0	66.0
	CI+	1.5	3.0	1.7	2.8	1.2		CI+	68.2	73.8	66.2	69.1	72.9
	CI-	0.0	0.0	0.0	0.1	0.0		CI-	59.2	59.1	55.5	54.5	58.6
Cocaine powder	%	4.7	6.2	3.3	6.9	1.9							
	CI+	7.5	11.4	6.1	11.6	5.0							
	CI-	2.8	3.0	1.5	3.7	0.5							
Amphetamines	%	5.1	6.7	3.7	8.2	1.2							
	CI+	7.6	11.0	7.2	12.8	3.5							
	CI-	3.3	3.6	1.6	4.8	0.3							
Ecstasy	%	5.9	6.8	5.1	10.2	0.3							
	CI+	8.6	11.3	8.3	15.1	1.8							
	CI-	3.8	3.6	2.8	6.5	0.0							

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**South Western Area Health Board (SWAHB)**

Table 14.2: SWAHB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	7.5	11.0	4.2
	CI+	11.1	16.8	7.1
	CI-	4.8	6.6	2.3
Cannabis	%	7.3	10.6	4.2
	CI+	10.9	16.5	7.1
	CI-	4.6	6.2	2.3
Heroin	%	0.4	0.5	0.3
	CI+	1.5	3.0	1.8
	CI-	0.0	0.0	0.0
Methadone	%	1.0	1.1	0.9
	CI+	2.3	3.8	2.7
	CI-	0.4	0.2	0.2
Other opiates <sup>2</sup>	%	1.5	1.3	1.7
	CI+	3.4	4.1	3.7
	CI-	0.5	0.2	0.6
Cocaine (total including crack)	%	1.5	2.1	0.8
	CI+	3.5	6.2	2.5
	CI-	0.5	0.4	0.2
Crack	%	0.2	0.0	0.3
	CI+	1.0	1.4	1.7
	CI-	0.0	0.0	0.0
Cocaine powder	%	1.5	2.1	0.8
	CI+	3.5	6.2	2.5
	CI-	0.5	0.4	0.2
Amphetamines	%	0.6	0.9	0.3
	CI+	1.8	3.6	1.8
	CI-	0.1	0.1	0.0
Ecstasy	%	1.3	1.3	1.3
	CI+	3.0	4.1	3.4
	CI-	0.4	0.2	0.4

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.3	0.5	0.2
	CI+	1.4	3.1	1.5
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.2	0.0	0.3
	CI+	1.0	1.4	1.8
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	6.2	4.6	7.7
	CI+	8.8	8.6	11.5
	CI-	4.1	2.1	4.8
Alcohol	%	86.0	88.0	84.0
	CI+	89.0	92.6	87.8
	CI-	82.5	82.0	79.7
Tobacco	%	43.8	47.6	40.1
	CI+	49.1	55.2	46.5
	CI-	38.6	40.2	33.9

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**South Western Area Health Board (SWAHB)**

**Table 14.3: SWAHB – Last Month Prevalence (%)**

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	4.3	6.9	1.8
	CI+	7.6	12.5	3.9
	CI-	2.2	3.3	0.7
Cannabis	%	3.9	6.1	1.8
	CI+	7.0	11.3	3.9
	CI-	1.9	2.8	0.7
Heroin	%	0.2	0.5	0.0
	CI+	1.4	3.0	1.1
	CI-	0.0	0.0	0.0
Methadone	%	0.6	0.3	0.9
	CI+	1.6	1.9	2.7
	CI-	0.2	0.0	0.2
Other opiates <sup>2</sup>	%	0.6	0.4	0.8
	CI+	1.6	2.7	2.4
	CI-	0.1	0.0	0.1
Cocaine (total including crack)	%	0.6	1.3	0.0
	CI+	1.9	4.2	1.1
	CI-	0.1	0.2	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.6	1.3	0.0
	CI+	1.9	4.2	1.1
	CI-	0.1	0.2	0.0
Amphetamines	%	0.2	0.5	0.0
	CI+	1.4	3.0	1.1
	CI-	0.0	0.0	0.0
Ecstasy	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.2	0.5	0.0
	CI+	1.4	3.1	1.1
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.0	0.0	0.0
	CI+	0.6	1.4	1.1
	CI-	0.0	0.0	0.0
Sedative, tranquillisers, anti-depressants	%	4.2	3.0	5.4
	CI+	6.7	6.6	9.1
	CI-	2.4	1.0	2.8
Alcohol	%	73.5	77.1	70.0
	CI+	77.9	83.6	75.5
	CI-	68.7	69.5	64.1
Tobacco	%	39.4	42.1	36.8
	CI+	44.2	48.4	43.4
	CI-	34.7	35.9	30.7

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 15 – Ireland

Southern Health Board (SHB)

Table 15.1: SHB – Lifetime Prevalence (%)

	All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		All Adults 15-64	Males	Females	Young Adults 15-34	Older Adults 35-64		
Any illegal drugs <sup>1</sup>	%	12.1	14.2	10.0	18.2	7.0	LSD	%	1.2	1.2	1.1	1.8	0.6
	CI+	15.8	20.5	13.7	24.8	10.4		CI+	2.3	3.3	2.7	4.2	2.2
	CI-	9.1	9.2	7.1	12.7	4.5		CI-	0.5	0.3	0.3	0.6	0.1
Cannabis	%	11.6	13.6	9.5	17.1	6.9	Magic mushrooms	%	2.9	3.1	2.6	4.6	1.4
	CI+	15.2	19.9	13.1	23.9	10.2		CI+	4.5	5.9	4.7	8.4	3.0
	CI-	8.5	8.7	6.6	11.7	4.4		CI-	1.7	1.4	1.3	2.2	0.5
Heroin	%	0.0	0.0	0.0	0.0	0.0	Solvents	%	1.1	1.6	0.6	2.2	0.2
	CI+	0.5	1.1	0.9	1.2	0.8		CI+	2.2	4.0	1.9	4.8	1.2
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	0.5	0.5	0.1	0.8	0.0
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%	1.3	1.9	0.7	2.9	0.0
	CI+	0.5	1.1	0.9	1.2	0.8		CI+	2.7	4.2	2.4	6.0	0.8
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	0.5	0.7	0.1	1.2	0.0
Other opiates <sup>2</sup>	%	0.8	0.6	1.0	1.2	0.4	Sedatives, tranquillisers, anti-depressants	%	9.9	5.5	14.3	6.3	12.9
	CI+	1.7	2.1	2.5	3.2	1.6		CI+	12.8	8.6	18.8	10.9	17.2
	CI-	0.3	0.1	0.3	0.3	0.1		CI-	7.4	3.3	10.6	3.2	9.3
Cocaine (total including crack)	%	1.9	2.2	1.5	2.8	1.0	Alcohol	%	91.2	93.3	89.1	93.2	89.5
	CI+	3.3	5.1	3.3	6.0	2.6		CI+	93.2	96.0	92.0	96.0	92.3
	CI-	0.9	0.7	0.6	1.1	0.3		CI-	88.8	89.6	85.6	89.5	86.2
Crack	%	0.0	0.0	0.0	0.0	0.0	Tobacco	%	54.7	53.2	56.2	51.9	57.0
	CI+	0.5	1.1	0.9	1.2	0.8		CI+	59.7	60.2	62.5	58.3	62.8
	CI-	0.0	0.0	0.0	0.0	0.0		CI-	49.7	46.2	49.8	45.5	51.1
Cocaine powder	%	1.9	2.2	1.5	2.8	1.0							
	CI+	3.3	5.1	3.3	6.0	2.6							
	CI-	0.9	0.7	0.6	1.1	0.3							
Amphetamines	%	1.8	2.3	1.3	3.0	0.8							
	CI+	3.5	6.0	2.9	6.9	2.3							
	CI-	0.8	0.6	0.4	1.0	0.2							
Ecstasy	%	2.8	3.9	1.7	5.6	0.4							
	CI+	5.0	7.8	3.5	10.4	1.6							
	CI-	1.4	1.6	0.7	2.6	0.0							

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Southern Health Board (SHB)**

Table 15.2: SHB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	4.7	5.5	4.0
	CI+	6.8	9.1	6.6
	CI-	3.1	3.0	2.2
Cannabis	%	4.4	4.8	4.0
	CI+	6.4	8.2	6.6
	CI-	2.8	2.5	2.2
Heroin	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.3	0.6	0.0
	CI+	1.0	2.1	0.9
	CI-	0.0	0.1	0.0
Cocaine (total including crack)	%	0.7	1.0	0.4
	CI+	1.8	3.2	1.8
	CI-	0.2	0.2	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.7	1.0	0.4
	CI+	1.8	3.2	1.8
	CI-	0.2	0.2	0.0
Amphetamines	%	0.4	0.8	0.0
	CI+	1.4	2.9	0.9
	CI-	0.0	0.1	0.0
Ecstasy	%	0.9	1.4	0.4
	CI+	2.4	4.7	1.8
	CI-	0.2	0.2	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.3	0.7	0.0
	CI+	1.2	2.6	0.9
	CI-	0.0	0.1	0.0
Magic mushrooms	%	0.3	0.3	0.4
	CI+	1.8	1.7	1.8
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.1	0.3	0.0
	CI+	0.8	1.7	0.9
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	4.1	2.7	5.7
	CI+	6.3	5.3	8.4
	CI-	2.6	1.1	3.6
Alcohol	%	85.1	88.0	82.1
	CI+	87.6	91.3	85.7
	CI-	82.3	84.0	78.1
Tobacco	%	34.7	32.0	37.5
	CI+	38.5	37.8	42.4
	CI-	31.1	26.6	32.8
				33.3
				26.0

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Southern Health Board (SHB)**

Table 15.3: SHB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	2.1	1.9	2.3
	CI+	3.8	4.3	4.7
	CI-	1.1	0.7	1.0
Cannabis	%	2.1	1.9	2.3
	CI+	3.8	4.3	4.7
	CI-	1.1	0.7	1.0
Heroin	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.1	0.3	0.0
	CI+	0.8	1.7	0.9
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.4	0.8	0.0
	CI+	1.4	2.9	0.9
	CI-	0.0	0.1	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.4	0.8	0.0
	CI+	1.4	2.9	0.9
	CI-	0.0	0.1	0.0
Amphetamines	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Ecstasy	%	0.2	0.4	0.0
	CI+	1.1	2.3	0.9
	CI-	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.1	0.3	0.0
	CI+	0.8	1.7	0.9
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.0	0.0	0.0
	CI+	0.5	1.1	0.9
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	3.2	2.7	3.9
	CI+	5.1	5.3	6.4
	CI-	1.9	1.1	2.1
Alcohol	%	75.3	79.0	71.6
	CI+	78.4	83.3	75.9
	CI-	72.1	74.2	66.9
Tobacco	%	31.4	28.3	34.6
	CI+	35.5	34.4	39.4
	CI-	27.5	22.7	30.0

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

## Prevalence Table 16 – Ireland

Western Health Board (WHB)

Table 16.1: WHB – Lifetime Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males			15-64	Males		
Any illegal drugs <sup>1</sup>	%	12.5	16.3	8.6	15.6	9.9	LSD	%
	CI+	18.3	25.3	14.4	23.1	16.9		CI+
	CI-	8.0	9.4	4.6	9.7	5.1		CI-
Cannabis	%	12.0	15.8	8.1	14.6	9.9	Magic mushrooms	%
	CI+	17.6	24.9	13.8	21.1	16.9		CI+
	CI-	7.7	9.1	4.2	9.4	5.1		CI-
Heroin	%	0.0	0.0	0.0	0.0	0.0	Solvents	%
	CI+	0.8	1.8	1.3	1.8	1.3		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Methadone	%	0.0	0.0	0.0	0.0	0.0	Poppers <sup>3</sup>	%
	CI+	0.8	1.8	1.3	1.8	1.3		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Other opiates <sup>2</sup>	%	1.1	0.4	1.8	0.9	1.3	Sedatives, tranquillisers, anti-depressants	%
	CI+	2.5	2.6	4.3	3.4	3.4		CI+
	CI-	0.4	0.0	0.6	0.1	0.3		CI-
Cocaine (total including crack)	%	1.6	2.1	1.2	2.2	1.2	Alcohol	%
	CI+	3.3	5.9	3.2	5.7	3.4		CI+
	CI-	0.7	0.5	0.3	0.6	0.3		CI-
Crack	%	0.1	0.0	0.3	0.3	0.0	Tobacco	%
	CI+	1.0	1.8	1.9	2.5	1.3		CI+
	CI-	0.0	0.0	0.0	0.0	0.0		CI-
Cocaine powder	%	1.5	2.1	0.9	1.9	1.2		
	CI+	3.2	5.9	2.8	5.4	3.4		
	CI-	0.6	0.5	0.1	0.4	0.3		
Amphetamines	%	1.4	2.2	0.5	2.0	0.9		
	CI+	3.2	5.4	2.6	6.0	3.1		
	CI-	0.4	0.6	0.0	0.3	0.1		
Ecstasy	%	1.8	2.2	1.4	2.5	1.2		
	CI+	3.6	5.4	3.6	6.6	3.4		
	CI-	0.7	0.6	0.3	0.6	0.3		

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Western Health Board (WHB)**

Table 16.2: WHB – Last Year Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	2.9	3.5	2.2
	CI+	5.4	7.3	5.4
	CI-	1.3	1.3	0.7
Cannabis	%	2.0	2.8	1.2
	CI+	4.0	6.1	3.3
	CI-	0.8	1.0	0.3
Heroin	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.4	0.0	0.9
	CI+	1.5	1.8	2.9
	CI-	0.1	0.0	0.1
Cocaine (total including crack)	%	0.7	1.3	0.0
	CI+	2.4	5.2	1.3
	CI-	0.1	0.1	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.7	1.3	0.0
	CI+	2.4	5.2	1.3
	CI-	0.1	0.1	0.0
Amphetamines	%	0.3	0.6	0.0
	CI+	1.6	3.6	1.3
	CI-	0.0	0.0	0.0
Ecstasy	%	0.3	0.0	0.5
	CI+	14	1.8	2.5
	CI-	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.3	0.0	0.6
	CI+	1.4	1.8	3.3
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	5.7	4.6	3.2
	CI+	8.1	8.4	7.0
	CI-	3.8	2.1	1.1
Alcohol	%	77.1	76.5	78.2
	CI+	80.8	82.2	83.7
	CI-	73.1	70.1	71.8
Tobacco	%	35.9	36.5	37.7
	CI+	40.9	45.6	46.1
	CI-	31.1	28.2	29.8

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

# Drug Use in Ireland & Northern Ireland

2002/2003 Drug Prevalence Survey: Health Boards (Ireland) and Health and Social Services Board (Northern Ireland) Updated Results:  
 Bulletin 2: Confidence Intervals (Incorporating the Design Effect)

**Updated Confidence Intervals**

**Western Health Board (WHB)**

Table 16.3: WHB – Last Month Prevalence (%)

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
Any illegal drugs <sup>1</sup>	%	1.9	2.1	1.7
	CI+	3.8	5.9	4.0
	CI-	0.8	0.5	0.5
Cannabis	%	1.3	1.4	1.2
	CI+	3.0	4.4	3.3
	CI-	0.4	0.2	0.3
Heroin	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Other opiates <sup>2</sup>	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Cocaine (total including crack)	%	0.4	0.7	0.0
	CI+	2.0	4.6	1.3
	CI-	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Cocaine powder	%	0.4	0.7	0.0
	CI+	2.0	4.6	1.3
	CI-	0.0	0.0	0.0
Amphetamines	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Ecstasy	%	0.3	0.0	0.5
	CI+	1.4	1.8	2.5
	CI-	0.0	0.0	0.0

	All Adults		Young Adults 15-34	Older Adults 35-64
	15-64	Males		
LSD	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Poppers <sup>3</sup>	%	0.0	0.0	0.0
	CI+	0.8	1.8	1.3
	CI-	0.0	0.0	0.0
Sedatives, tranquillisers, anti-depressants	%	3.8	2.8	4.9
	CI+	5.9	6.1	8.2
	CI-	2.3	1.0	2.7
Alcohol	%	67.0	70.2	63.7
	CI+	71.2	76.4	69.4
	CI-	62.7	63.5	57.8
Tobacco	%	31.5	30.5	32.5
	CI+	37.3	41.1	41.9
	CI-	26.0	21.2	24.0

<sup>1</sup> For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

<sup>2</sup> Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

<sup>3</sup> Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.



**DAIRU**  
Drug and Alcohol  
Information  
and Research Unit

Department of  
Health, Social Services  
and Public Safety  
[www.dhsspsni.gov.uk](http://www.dhsspsni.gov.uk)  
AN RIALT  
Sláinte, Seirbhísí Sóisialta  
agus Sébháilteachta Poiblí  
MINISTERSTEW  
Poblacht, Resydénter Heisín  
an Fowk Sicar

  
**Ipsos MORI**

 *Northern Ireland*  
**Statistics &  
Research** agency

  
**NDP**  
National Development Plan 2007 - 2013  
Transforming Ireland