

**Updated
Confidence
Intervals**

Drug Use in Ireland & Northern Ireland

Updated Results from the 2002/2003 Drug Prevalence Survey – *Bulletin 1*



Prevalence Tables – Ireland

Table 1.1: Ireland – Lifetime Prevalence (%)

	All Adults			Young Adults						
	15-64		Males	Females	15-34	15-24	25-34	35-44	45-54	55-64
	%									
Any illegal drugs ¹	%	18.5	24.0	13.1	26.0	24.9	27.1	17.7	10.6	4.2
	CI+	20.1	26.6	14.6	28.5	28.7	30.4	20.0	13.3	5.8
	CI-	17.0	21.5	11.6	23.5	21.3	23.9	15.5	8.3	2.9
Cannabis	%	17.4	22.4	12.3	24.0	22.8	25.2	17.3	10.1	3.6
	CI+	18.9	24.9	13.8	26.5	26.5	28.3	19.6	12.7	5.2
	CI-	15.9	20.0	10.9	21.6	19.4	22.2	15.1	7.9	2.4
Heroin	%	0.5	0.7	0.3	0.7	0.4	1.0	0.5	0.3	0.0
	CI+	0.8	1.2	0.6	1.3	1.2	2.0	1.0	1.0	0.5
	CI-	0.3	0.3	0.1	0.4	0.1	0.5	0.2	0.0	0.0
Methadone	%	0.3	0.4	0.3	0.6	0.2	1.0	0.3	0.0	0.0
	CI+	0.6	0.8	0.7	1.1	1.0	1.8	0.8	0.4	0.5
	CI-	0.2	0.1	0.1	0.3	0.0	0.4	0.1	0.0	0.0
Other opiates ²	%	3.0	2.0	4.0	3.0	1.5	4.5	3.8	3.0	1.9
	CI+	3.7	2.8	5.0	4.1	2.8	6.1	5.0	4.7	3.2
	CI-	2.4	1.4	3.1	2.1	0.7	3.2	2.8	1.8	1.1
Cocaine (total including crack)	%	3.0	4.3	1.6	4.7	5.1	4.2	2.8	0.9	0.0
	CI+	3.6	5.4	2.3	5.9	7.3	5.8	3.9	1.8	0.5
	CI-	2.4	3.4	1.1	3.6	3.4	3.0	2.0	0.3	0.0
Crack	%	0.3	0.5	0.1	0.5	0.5	0.5	0.4	0.0	0.0
	CI+	0.5	1.0	0.3	0.9	1.4	1.1	1.0	0.4	0.5
	CI-	0.2	0.2	0.0	0.2	0.1	0.1	0.2	0.0	0.0
Cocaine powder	%	2.9	4.1	1.6	4.5	4.9	4.2	2.6	0.9	0.0
	CI+	3.5	5.2	2.2	5.8	7.0	5.7	3.7	1.8	0.5
	CI-	2.3	3.2	1.1	3.5	3.3	2.9	1.8	0.3	0.0
Amphetamines	%	3.0	4.0	1.9	4.8	4.4	5.3	2.2	0.8	0.4
	CI+	3.6	5.1	2.6	6.1	6.4	6.9	3.2	1.8	1.2
	CI-	2.4	3.1	1.3	3.7	2.8	3.9	1.4	0.3	0.1

	All Adults			Young Adults						
	15-64		Males	Females	15-34	15-24	25-34	35-44	45-54	55-64
	%									
Ecstasy	%	3.7	4.9	2.6	7.1	7.7	6.4	1.6	0.1	0.0
	CI+	4.4	6.1	3.3	8.6	10.2	8.3	2.5	0.6	0.5
	CI-	3.1	3.8	1.9	5.8	5.7	4.9	1.0	0.0	0.0
LSD	%	2.9	4.4	1.4	4.6	3.9	5.3	2.2	1.2	0.3
	CI+	3.5	5.5	2.0	5.7	5.7	7.0	3.2	2.3	1.1
	CI-	2.4	3.4	1.0	3.6	2.5	3.9	1.5	0.5	0.1
Magic mushrooms	%	3.9	5.7	2.0	5.9	5.5	6.3	4.2	0.8	0.3
	CI+	4.5	6.9	2.7	7.3	7.6	8.2	5.5	1.6	1.0
	CI-	3.3	4.7	1.5	4.7	3.9	4.7	3.2	0.3	0.0
Solvents	%	1.7	2.2	1.1	3.3	3.6	3.1	0.4	0.0	0.2
	CI+	2.2	3.1	1.7	4.4	5.3	4.4	0.9	0.4	0.9
	CI-	1.3	1.5	0.7	2.4	2.2	2.1	0.1	0.0	0.0
Poppers ³	%	2.6	3.9	1.3	4.7	4.7	4.8	1.3	0.4	0.1
	CI+	3.2	5.0	1.9	6.1	6.7	6.5	2.1	1.1	0.7
	CI-	2.1	2.9	0.9	3.6	3.1	3.3	0.7	0.1	0.0
Sedatives, Tranquillisers, Anti-depressants	%	12.1	9.3	15.0	8.2	6.5	9.9	12.2	15.3	21.8
	CI+	13.2	10.6	16.5	9.6	8.6	12.0	14.2	17.9	24.9
	CI-	11.2	8.0	13.7	6.9	4.7	8.1	10.4	13.0	19.0

¹ For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

² Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, naps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

³ Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

Prevalence Tables – Ireland

Table 1.2: Ireland – Last Year Prevalence (%)

	All Adults		Young Adults							
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Any illegal drugs ¹	%	5.6	7.8	3.4	9.7	12.7	6.6	3.4	1.1	0.4
	CI+	6.5	9.3	4.2	11.5	15.8	8.5	4.6	2.2	1.1
	CI-	4.8	6.5	2.6	8.1	10.0	5.0	2.4	0.5	0.1
Cannabis	%	5.0	7.2	2.9	8.6	11.0	6.2	3.3	1.1	0.4
	CI+	5.9	8.6	3.7	10.3	13.9	8.0	4.5	2.2	1.1
	CI-	4.3	5.9	2.2	7.1	8.4	4.6	2.4	0.5	0.1
Heroin	%	0.1	0.1	0.1	0.2	0.2	0.2	0.0	0.0	0.0
	CI+	0.3	0.3	0.4	0.6	1.0	0.7	0.3	0.4	0.5
	CI-	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0
Methadone	%	0.2	0.2	0.2	0.4	0.2	0.5	0.2	0.0	0.0
	CI+	0.4	0.5	0.5	0.8	1.0	1.2	0.6	0.4	0.5
	CI-	0.1	0.0	0.1	0.1	0.0	0.2	0.0	0.0	0.0
Other opiates ²	%	0.5	0.4	0.5	0.6	0.6	0.5	0.3	0.5	0.4
	CI+	0.8	0.9	0.9	1.1	1.5	1.2	0.8	1.2	1.2
	CI-	0.3	0.2	0.3	0.2	0.2	0.2	0.1	0.1	0.1
Cocaine (total including crack)	%	1.1	1.7	0.5	2.0	2.7	1.3	0.5	0.2	0.0
	CI+	1.5	2.5	0.8	2.9	4.4	2.3	1.1	0.8	0.5
	CI-	0.7	1.1	0.2	1.3	1.5	0.6	0.2	0.0	0.0
Crack	%	0.1	0.1	0.1	0.2	0.1	0.2	0.0	0.0	0.0
	CI+	0.2	0.3	0.3	0.5	0.8	0.7	0.3	0.4	0.5
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cocaine powder	%	1.1	1.7	0.5	2.0	2.7	1.3	0.5	0.2	0.0
	CI+	1.5	2.5	0.8	2.9	4.4	2.3	1.1	0.8	0.5
	CI-	0.7	1.1	0.2	1.3	1.5	0.6	0.2	0.0	0.0
Amphetamines	%	0.4	0.7	0.2	0.8	1.2	0.3	0.2	0.0	0.0
	CI+	0.7	1.2	0.4	1.4	2.3	1.0	0.7	0.4	0.5
	CI-	0.2	0.3	0.1	0.4	0.5	0.1	0.0	0.0	0.0

	All Adults		Young Adults							
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Ecstasy	%	1.1	1.5	0.6	2.3	2.8	1.7	0.0	0.0	0.0
	CI+	1.5	2.3	1.0	3.1	4.4	2.8	0.3	0.4	0.5
	CI-	0.8	1.0	0.3	1.6	1.7	0.9	0.0	0.0	0.0
LSD	%	0.1	0.2	0.0	0.2	0.2	0.1	0.0	0.0	0.0
	CI+	0.2	0.5	0.1	0.5	0.9	0.5	0.3	0.4	0.5
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Magic mushrooms	%	0.4	0.6	0.1	0.7	1.1	0.3	0.2	0.0	0.1
	CI+	0.6	1.2	0.3	1.3	2.1	0.8	0.6	0.4	0.7
	CI-	0.2	0.3	0.0	0.3	0.4	0.1	0.0	0.0	0.0
Solvents	%	0.1	0.0	0.1	0.1	0.2	0.1	0.0	0.0	0.0
	CI+	0.2	0.3	0.3	0.4	0.8	0.5	0.3	0.4	0.5
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Poppers ³	%	0.4	0.4	0.4	0.8	0.9	0.6	0.1	0.0	0.0
	CI+	0.6	0.9	0.7	1.3	1.9	1.3	0.5	0.4	0.5
	CI-	0.2	0.2	0.2	0.4	0.4	0.2	0.0	0.0	0.0
Sedatives, Tranquillisers	%	5.7	4.2	7.3	3.7	3.6	3.9	4.5	7.7	12.1
	CI+	6.5	5.2	8.3	4.7	5.2	5.3	5.8	9.7	14.7
	CI-	5.0	3.3	6.3	2.9	2.4	2.7	3.4	6.0	9.9

¹ For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

² Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

³ Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

Prevalence Tables – Ireland

Table 1.3: Ireland – Last Month Prevalence (%)

	All Adults			Young Adults						
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Any illegal drugs ¹	%	3.0	4.1	1.7	5.2	6.9	3.4	1.6	0.6	0.3
	CI+	3.6	5.3	2.4	6.6	9.4	4.8	2.5	1.5	1.0
	CI-	2.4	3.1	1.2	4.0	4.9	2.2	1.0	0.1	0.0
Cannabis	%	2.6	3.4	1.7	4.3	5.6	3.0	1.6	0.6	0.3
	CI+	3.2	4.5	2.3	5.6	8.0	4.4	2.5	1.5	1.0
	CI-	2.0	2.5	1.2	3.3	3.7	1.9	1.0	0.1	0.0
Heroin	%	0.1	0.1	0.1	0.1	0.2	0.0	0.0	0.0	0.0
	CI+	0.2	0.3	0.3	0.5	1.0	0.3	0.3	0.4	0.5
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Methadone	%	0.1	0.1	0.2	0.2	0.1	0.3	0.2	0.0	0.0
	CI+	0.3	0.3	0.5	0.5	0.7	0.9	0.6	0.4	0.5
	CI-	0.1	0.0	0.1	0.0	0.0	0.1	0.0	0.0	0.0
Other opiates ²	%	0.2	0.2	0.1	0.1	0.1	0.2	0.1	0.3	0.1
	CI+	0.3	0.5	0.4	0.5	0.8	0.6	0.4	0.9	0.7
	CI-	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cocaine (total including crack)	%	0.3	0.7	0.0	0.7	0.9	0.4	0.1	0.0	0.0
	CI+	0.6	1.2	0.1	1.2	2.0	1.1	0.5	0.4	0.5
	CI-	0.2	0.3	0.0	0.3	0.3	0.1	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	CI+	0.1	0.2	0.1	0.2	0.4	0.3	0.3	0.4	0.5
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cocaine powder	%	0.3	0.7	0.0	0.7	0.9	0.4	0.1	0.0	0.0
	CI+	0.6	1.2	0.1	1.2	2.0	1.1	0.5	0.4	0.5
	CI-	0.2	0.3	0.0	0.3	0.3	0.1	0.0	0.0	0.0
Amphetamines	%	0.2	0.2	0.1	0.3	0.6	0.0	0.1	0.0	0.0
	CI+	0.3	0.6	0.3	0.7	1.5	0.3	0.5	0.4	0.5
	CI-	0.1	0.1	0.0	0.1	0.1	0.0	0.0	0.0	0.0

	All Adults			Young Adults					
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64
Ecstasy	%	0.3	0.6	0.0	0.6	0.8	0.5	0.0	0.0
	CI+	0.5	1.0	0.2	1.2	1.8	1.2	0.3	0.4
	CI-	0.2	0.3	0.0	0.3	0.3	0.1	0.0	0.0
LSD	%	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0
	CI+	0.1	0.3	0.1	0.3	0.4	0.5	0.3	0.4
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Magic mushrooms	%	0.1	0.1	0.0	0.1	0.1	0.0	0.0	0.1
	CI+	0.2	0.4	0.1	0.4	0.8	0.3	0.3	0.4
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Solvents	%	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0
	CI+	0.1	0.3	0.1	0.3	0.7	0.3	0.3	0.4
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Poppers ³	%	0.1	0.1	0.0	0.2	0.3	0.0	0.0	0.0
	CI+	0.2	0.4	0.2	0.5	1.1	0.3	0.3	0.4
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Sedatives, Tranquillisers, Anti-depressants	%	4.0	3.0	4.9	2.1	2.0	2.2	2.6	6.2
	CI+	4.6	4.0	5.8	2.9	3.3	3.4	3.6	8.0
	CI-	3.4	2.3	4.1	1.5	1.1	1.4	1.8	4.6
									9.7
									12.0

¹ For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

² Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, naps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

³ Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

Prevalence Tables – Northern Ireland

Table 2.1: Northern Ireland – Lifetime Prevalence (%)

	All Adults		Young Adults							
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Any illegal drugs ¹	%	20.0	26.7	13.5	30.9	28.5	33.2	18.4	8.9	4.0
	CI+	21.8	29.6	15.4	34.2	33.8	37.0	21.4	11.5	5.8
	CI-	18.3	23.9	11.7	27.8	23.7	29.6	15.7	6.8	2.7
Cannabis	%	16.8	23.3	10.5	25.1	24.0	26.2	16.9	8.0	3.1
	CI+	18.5	26.0	12.1	28.2	29.4	29.4	19.8	10.5	4.7
	CI-	15.3	20.6	9.0	22.3	19.2	23.1	14.3	6.0	1.9
Heroin	%	0.2	0.4	0.0	0.3	0.4	0.3	0.3	0.0	0.0
	CI+	0.4	0.9	0.2	0.9	1.6	0.9	1.0	0.6	0.6
	CI-	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0
Methadone	%	0.1	0.2	0.1	0.2	0.2	0.1	0.0	0.2	0.1
	CI+	0.3	0.6	0.4	0.7	1.4	0.7	0.4	0.9	0.8
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other opiates ²	%	18.0	16.4	19.5	17.9	13.0	22.7	22.8	15.2	14.3
	CI+	20.2	19.2	22.1	20.9	17.6	26.1	26.5	18.8	17.2
	CI-	15.9	13.9	17.1	15.1	9.1	19.6	19.4	12.0	11.6
Cocaine (total including crack)	%	1.6	2.8	0.5	2.9	2.6	3.2	1.5	0.1	0.0
	CI+	2.2	3.9	1.0	4.1	4.8	4.6	2.6	0.8	0.6
	CI-	1.2	1.9	0.3	2.0	1.2	2.1	0.8	0.0	0.0
Crack	%	0.2	0.4	0.1	0.4	0.6	0.3	0.3	0.0	0.0
	CI+	0.5	1.0	0.3	1.0	1.9	0.9	0.9	0.6	0.6
	CI-	0.1	0.2	0.0	0.1	0.1	0.0	0.0	0.0	0.0
Cocaine powder	%	1.6	2.7	0.5	2.9	2.6	3.1	1.4	0.1	0.0
	CI+	2.1	3.8	0.9	4.0	4.8	4.5	2.4	0.8	0.6
	CI-	1.2	1.8	0.2	2.0	1.2	2.0	0.7	0.0	0.0
Amphetamines	%	4.0	6.3	1.8	7.2	6.5	8.0	2.6	0.8	0.8
	CI+	4.9	7.8	2.6	9.0	9.8	10.1	3.9	1.8	1.8
	CI-	3.3	5.0	1.2	5.7	4.0	6.1	1.6	0.2	0.3

	All Adults		Young Adults							
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Ecstasy	%	5.8	8.3	3.3	11.3	10.2	12.4	2.9	0.6	0.0
	CI+	6.7	10.0	4.3	13.5	13.9	14.9	4.4	1.6	0.6
	CI-	4.9	6.8	2.4	9.4	7.2	10.2	1.8	0.2	0.0
LSD	%	4.5	7.0	2.0	7.8	5.9	9.6	3.7	1.0	0.4
	CI+	5.4	8.7	2.8	9.5	8.9	11.7	5.5	2.3	1.3
	CI-	3.7	5.6	1.4	6.2	3.7	7.6	2.4	0.3	0.1
Magic mushrooms	%	4.3	6.3	2.4	6.4	4.9	7.9	5.9	0.7	0.3
	CI+	5.2	7.9	3.2	8.0	7.5	9.9	7.9	1.7	1.1
	CI-	3.6	4.9	1.8	5.0	2.9	6.2	4.3	0.2	0.0
Solvents	%	2.9	4.1	1.7	5.4	4.9	5.8	2.0	0.2	0.3
	CI+	3.6	5.5	2.4	6.9	7.6	7.6	3.3	0.9	1.1
	CI-	2.3	3.0	1.2	4.1	3.0	4.3	1.2	0.0	0.0
Poppers ³	%	5.5	7.3	3.8	10.9	10.6	11.1	2.6	0.8	0.0
	CI+	6.6	9.0	5.0	13.3	14.9	13.5	4.1	1.9	0.6
	CI-	4.5	5.9	2.7	8.8	7.2	9.1	1.6	0.3	0.0
Sedatives, Tranquillisers, Anti-depressants	%	22.1	15.5	28.5	16.2	10.9	21.4	24.6	27.2	29.4
	CI+	23.8	17.7	30.7	18.7	14.5	24.5	27.8	30.9	33.0
	CI-	20.4	13.4	26.5	13.8	7.7	18.5	21.5	23.7	25.9

¹ For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

² Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, naps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

³ Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

Prevalence Tables – Northern Ireland

Table 2.2: Northern Ireland – Last Year Prevalence (%)

	All Adults		Young Adults							
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Any illegal drugs ¹	%	6.4	9.7	3.1	11.6	14.5	8.8	3.7	1.7	0.6
	CI+	7.4	11.6	4.1	13.9	18.8	11.0	5.2	3.1	1.5
	CI-	5.4	8.0	2.3	9.7	10.9	7.0	2.5	0.9	0.2
Cannabis	%	5.4	8.7	2.2	9.8	12.1	7.4	3.7	1.4	0.1
	CI+	6.4	10.5	3.1	11.9	16.4	9.4	5.2	2.6	0.8
	CI-	4.5	7.0	1.4	7.9	8.7	5.8	2.5	0.6	0.0
Heroin	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	CI+	0.1	0.2	0.2	0.3	0.7	0.4	0.4	0.6	0.6
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Methadone	%	0.0	0.0	0.1	0.0	0.0	0.0	0.2	0.0	0.0
	CI+	0.2	0.2	0.3	0.3	0.7	0.4	0.4	0.9	0.6
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other opiates ²	%	8.0	6.9	9.0	7.5	4.1	10.9	10.8	6.0	7.3
	CI+	9.3	8.5	10.6	9.4	6.7	13.4	13.5	8.5	9.5
	CI-	6.8	5.5	7.6	6.0	2.3	8.7	8.5	4.0	5.4
Cocaine (total including crack)	%	0.5	1.0	0.1	1.0	1.4	0.7	0.3	0.0	0.0
	CI+	0.9	1.8	0.4	1.8	3.2	1.5	1.0	0.6	0.6
	CI-	0.3	0.4	0.0	0.5	0.4	0.2	0.0	0.0	0.0
Crack	%	0.0	0.1	0.0	0.1	0.2	0.0	0.0	0.0	0.0
	CI+	0.2	0.5	0.2	0.5	1.2	0.4	0.4	0.6	0.6
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cocaine powder	%	0.5	0.9	0.1	0.9	1.2	0.7	0.3	0.0	0.0
	CI+	0.8	1.6	0.4	1.7	2.9	1.5	1.0	0.6	0.6
	CI-	0.2	0.4	0.0	0.4	0.3	0.2	0.0	0.0	0.0
Amphetamines	%	0.8	1.4	0.2	1.5	2.0	1.0	0.3	0.2	0.3
	CI+	1.3	2.3	0.5	2.5	4.2	2.0	1.0	0.9	1.0
	CI-	0.5	0.8	0.0	0.8	0.7	0.5	0.0	0.0	0.0

	All Adults		Young Adults							
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Ecstasy	%	1.6	2.7	0.6	3.3	4.8	1.7	0.8	0.0	0.0
	CI+	2.2	3.9	1.1	4.5	7.6	2.8	1.7	0.9	0.6
	CI-	1.2	1.8	0.3	2.3	2.8	1.0	0.3	0.0	0.0
LSD	%	0.1	0.0	0.1	0.1	0.2	0.1	0.0	0.0	0.0
	CI+	0.2	0.2	0.4	0.5	1.1	0.6	0.4	0.6	0.6
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Magic mushrooms	%	0.2	0.4	0.1	0.3	0.5	0.1	0.1	0.2	0.0
	CI+	0.5	0.9	0.3	0.9	1.8	0.7	0.8	0.9	0.6
	CI-	0.1	0.1	0.0	0.1	0.1	0.0	0.0	0.0	0.0
Solvents	%	0.2	0.3	0.1	0.3	0.7	0.0	0.0	0.0	0.2
	CI+	0.4	0.8	0.3	0.9	2.1	0.4	0.4	0.6	0.9
	CI-	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0
Poppers ³	%	0.5	0.7	0.3	1.1	1.8	0.4	0.0	0.0	0.0
	CI+	0.8	1.4	0.6	1.9	3.7	1.0	0.4	0.6	0.6
	CI-	0.2	0.3	0.1	0.5	0.7	0.1	0.0	0.0	0.0
Sedatives, Tranquillisers, Anti-depressants	%	12.6	8.7	16.5	8.6	4.8	12.3	14.3	17.2	16.4
	CI+	13.9	10.3	18.3	10.3	7.3	14.7	17.0	20.4	19.5
	CI-	11.4	7.3	14.8	7.0	2.9	10.2	11.9	14.3	13.7

¹ For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

² Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, naps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

³ Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

Prevalence Tables – Northern Ireland

Table 2.3: Northern Ireland – Last Month Prevalence (%)

	All Adults			Young Adults						
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Any illegal drugs ¹	%	3.4	5.7	1.1	6.1	9.0	3.3	2.0	0.7	0.5
	CI+	4.2	7.3	1.7	7.9	12.7	4.8	3.2	1.7	1.3
	CI-	2.6	4.3	0.6	4.6	6.0	2.2	1.2	0.2	0.1
Cannabis	%	2.9	5.1	0.8	5.4	7.9	3.0	2.0	0.5	0.0
	CI+	3.7	6.7	1.3	7.1	11.5	4.4	3.2	1.4	0.6
	CI-	2.3	3.8	0.4	4.0	5.1	2.0	1.2	0.1	0.0
Heroin	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	CI+	0.1	0.2	0.2	0.3	0.7	0.4	0.4	0.6	
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Methadone	%	0.0	0.0	0.1	0.0	0.0	0.0	0.2	0.0	
	CI+	0.2	0.2	0.3	0.3	0.7	0.4	0.4	0.6	
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Other opiates ²	%	4.1	3.1	5.2	3.3	2.0	4.5	5.7	4.5	3.8
	CI+	4.9	4.0	6.3	4.4	3.7	6.1	7.6	6.4	5.6
	CI-	3.4	2.3	4.2	2.3	0.9	3.2	4.2	3.0	2.5
Cocaine (total including crack)	%	0.1	0.2	0.0	0.2	0.2	0.1	0.0	0.0	
	CI+	0.3	0.6	0.2	0.7	1.5	0.7	0.4	0.6	
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Crack	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	CI+	0.1	0.2	0.2	0.3	0.7	0.4	0.4	0.6	
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Cocaine powder	%	0.1	0.2	0.0	0.2	0.2	0.1	0.0	0.0	
	CI+	0.3	0.6	0.2	0.7	1.5	0.7	0.4	0.6	
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Amphetamines	%	0.2	0.5	0.0	0.3	0.4	0.3	0.0	0.2	0.3
	CI+	0.5	1.0	0.2	0.9	1.7	0.9	0.4	0.9	1.0
	CI-	0.1	0.2	0.0	0.1	0.0	0.0	0.0	0.0	0.0

	All Adults			Young Adults						
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Ecstasy	%	0.5	0.9	0.1	1.1	2.0	0.3	0.1	0.0	0.0
	CI+	0.9	1.6	0.5	1.9	3.8	0.9	0.7	0.6	0.6
	CI-	0.3	0.4	0.0	0.6	0.8	0.0	0.0	0.0	0.0
LSD	%	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
	CI+	0.1	0.2	0.3	0.4	0.7	0.6	0.4	0.6	0.6
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.0
	CI+	0.2	0.4	0.2	0.3	0.7	0.4	0.8	0.6	0.6
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Solvents	%	0.1	0.1	0.1	0.1	0.2	0.0	0.0	0.0	0.2
	CI+	0.2	0.3	0.3	0.5	1.1	0.4	0.4	0.6	0.9
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Poppers ³	%	0.2	0.2	0.1	0.3	0.6	0.1	0.0	0.0	0.0
	CI+	0.4	0.7	0.4	0.9	1.9	0.6	0.4	0.6	0.6
	CI-	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0
Sedatives, Tranquillisers, Anti-depressants	%	9.8	7.0	12.5	5.4	2.5	8.2	10.7	15.5	14.2
	CI+	10.9	8.5	14.1	6.8	4.5	10.2	13.2	18.7	17.1
	CI-	8.7	5.7	11.1	4.2	1.2	6.4	8.6	12.7	11.7

¹ For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

² Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

³ Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

Prevalence Tables – Island of Ireland

Table 3.1: Island of Ireland – Lifetime Prevalence (%)

	All Adults			Young Adults						
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Any illegal drugs ¹	%	19.0	24.8	13.2	27.4	26.0	28.9	17.9	10.1	4.1
	CI+	20.4	27.2	14.5	30.1	29.4	32.3	19.9	11.9	5.3
	CI-	18.0	23.2	12.1	26.0	23.3	27.3	16.2	8.1	3.1
Cannabis	%	17.2	22.6	11.8	24.3	23.2	25.5	17.2	9.5	3.5
	CI+	18.3	24.7	12.7	26.4	26.4	28.0	19.0	11.2	4.4
	CI-	16.0	20.9	10.5	22.6	20.4	23.3	15.4	7.6	2.5
Heroin	%	0.4	0.6	0.2	0.6	0.4	0.8	0.4	0.2	0.0
	CI+	0.6	0.9	0.4	0.9	1.0	1.3	0.8	0.7	0.3
	CI-	0.2	0.3	0.1	0.3	0.1	0.3	0.2	0.0	0.0
Methadone	%	0.3	0.3	0.3	0.5	0.2	0.7	0.2	0.0	0.0
	CI+	0.4	0.6	0.5	0.8	0.8	1.1	0.5	0.4	0.4
	CI-	0.1	0.1	0.1	0.2	0.0	0.3	0.0	0.0	0.0
Other opiates ²	%	7.4	6.2	8.5	7.3	4.9	9.8	9.3	6.5	5.5
	CI+	10.2	9.3	11.6	10.5	7.6	14.1	12.9	9.7	9.1
	CI-	8.4	7.1	9.3	7.9	4.4	10.9	9.9	6.5	6.3
Cocaine (total including crack)	%	2.6	3.9	1.3	4.2	4.4	3.9	2.4	0.6	0.0
	CI+	2.9	4.5	1.6	4.9	5.8	4.9	3.1	1.2	0.3
	CI-	2.0	3.0	0.8	3.2	2.9	2.9	1.6	0.2	0.0
Crack	%	0.3	0.5	0.1	0.5	0.5	0.4	0.4	0.0	0.0
	CI+	0.4	0.8	0.2	0.8	1.2	0.8	0.7	0.2	0.3
	CI-	0.2	0.3	0.0	0.2	0.2	0.1	0.2	0.0	0.0
Cocaine powder	%	2.5	3.7	1.3	4.1	4.3	3.9	2.3	0.6	0.0
	CI+	2.8	4.3	1.6	4.7	5.6	4.8	2.9	1.2	0.3
	CI-	1.9	2.8	0.8	3.1	2.8	2.8	1.5	0.2	0.0
Amphetamines	%	3.3	4.7	1.9	5.5	5.0	6.1	2.3	0.8	0.5
	CI+	3.9	5.9	2.4	6.9	6.9	7.8	3.1	1.5	1.2
	CI-	3.0	4.2	1.4	4.9	3.8	5.3	1.7	0.4	0.3

	All Adults			Young Adults						
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Ecstasy	%	4.3	5.9	2.8	8.3	8.5	8.2	2.0	0.3	0.0
	CI+	5.2	7.3	3.4	10.1	10.7	10.6	2.9	0.7	0.3
	CI-	4.0	5.4	2.3	7.7	6.9	7.7	1.5	0.1	0.0
LSD	%	3.4	5.1	1.6	5.5	4.5	6.5	2.7	1.1	0.4
	CI+	4.1	6.4	2.1	6.9	6.2	8.5	3.7	1.9	0.9
	CI-	3.1	4.7	1.3	5.0	3.4	6.0	2.1	0.6	0.1
Magic mushrooms	%	4.0	5.9	2.1	6.1	5.4	6.7	4.7	0.7	0.3
	CI+	4.6	6.9	2.7	7.1	6.9	8.4	5.9	1.3	0.7
	CI-	3.6	5.1	1.7	5.2	3.9	5.7	4.0	0.4	0.1
Solvents	%	2.0	2.8	1.3	3.9	4.0	3.9	0.8	0.0	0.3
	CI+	2.6	3.8	1.8	5.1	5.6	5.3	1.6	0.4	0.7
	CI-	1.8	2.4	1.0	3.4	2.9	3.4	0.6	0.0	0.1
Poppers ³	%	3.5	4.9	2.0	6.5	6.4	6.6	1.7	0.5	0.1
	CI+	4.4	6.3	2.9	8.5	8.8	9.0	2.5	1.1	0.4
	CI-	3.3	4.5	1.8	6.2	5.3	6.2	1.3	0.2	0.0
Sedatives, Tranquillisers, Anti-depressants	%	15.0	11.1	19.0	10.5	7.7	13.3	15.8	18.8	24.0
	CI+	17.2	13.1	21.9	12.7	10.0	16.7	18.8	22.4	27.7
	CI-	15.4	10.8	19.4	10.3	6.5	13.2	15.5	18.2	23.1

¹ For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

² Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, napps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

³ Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

Prevalence Tables – Island of Ireland

Table 3.2: Island of Ireland – Last Year Prevalence (%)

	All Adults			Young Adults						
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Any illegal drugs ¹	%	5.8	8.3	3.3	10.3	13.3	7.3	3.5	1.3	0.4
	CI+	6.6	9.8	3.9	11.9	16.0	9.0	4.5	2.2	1.0
	CI-	5.3	7.5	2.7	9.2	11.1	6.3	2.7	0.8	0.2
Cannabis	%	5.2	7.6	2.7	9.0	11.3	6.5	3.4	1.2	0.3
	CI+	5.8	8.9	3.2	10.4	13.9	8.1	4.4	2.0	0.7
	CI-	4.6	6.7	2.1	7.9	9.3	5.5	2.7	0.7	0.1
Heroin	%	0.1	0.0	0.1	0.2	0.2	0.1	0.0	0.0	0.0
	CI+	0.2	0.2	0.2	0.4	0.6	0.4	0.2	0.2	0.3
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Methadone	%	0.2	0.1	0.2	0.3	0.2	0.4	0.1	0.0	0.0
	CI+	0.3	0.3	0.4	0.5	0.6	0.7	0.4	0.4	0.3
	CI-	0.1	0.0	0.1	0.1	0.0	0.1	0.0	0.0	0.0
Other opiates ²	%	2.7	2.3	3.0	2.6	1.6	3.6	3.4	2.1	2.4
	CI+	4.1	3.8	4.6	4.2	2.9	6.1	5.5	3.7	4.7
	CI-	3.1	2.6	3.5	2.8	1.2	4.1	3.6	1.9	2.7
Cocaine (total including crack)	%	0.9	1.5	0.4	1.7	2.3	1.1	0.4	0.1	0.0
	CI+	1.2	2.0	0.6	2.2	3.4	1.7	0.8	0.5	0.3
	CI-	0.6	0.9	0.2	1.1	1.3	0.5	0.2	0.0	0.0
Crack	%	0.1	0.1	0.1	0.1	0.2	0.1	0.0	0.0	0.0
	CI+	0.2	0.3	0.2	0.4	0.6	0.4	0.2	0.2	0.3
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cocaine powder	%	0.9	1.5	0.4	1.7	2.3	1.1	0.4	0.1	0.0
	CI+	1.1	1.9	0.6	2.2	3.3	1.7	0.8	0.5	0.3
	CI-	0.6	0.9	0.2	1.1	1.2	0.5	0.2	0.0	0.0
Amphetamines	%	0.5	0.9	0.2	1.0	1.4	0.5	0.2	0.0	0.1
	CI+	0.8	1.4	0.3	1.6	2.5	1.1	0.6	0.4	0.5
	CI-	0.4	0.6	0.1	0.7	0.8	0.3	0.1	0.0	0.0

	All Adults			Young Adults						
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Ecstasy	%	1.2	1.9	0.6	2.5	3.4	1.7	0.2	0.0	0.0
	CI+	1.6	2.6	0.9	3.4	5.0	2.5	0.7	0.4	0.3
	CI-	1.0	1.5	0.4	2.1	2.5	1.1	0.1	0.0	0.0
LSD	%	0.1	0.1	0.0	0.2	0.2	0.1	0.0	0.0	0.0
	CI+	0.2	0.3	0.2	0.4	0.7	0.4	0.2	0.2	0.3
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Magic mushrooms	%	0.3	0.5	0.1	0.6	0.9	0.2	0.2	0.0	0.1
	CI+	0.5	0.9	0.2	1.0	1.7	0.6	0.4	0.4	0.4
	CI-	0.2	0.3	0.0	0.3	0.4	0.0	0.0	0.0	0.0
Solvents	%	0.1	0.1	0.1	0.2	0.3	0.1	0.0	0.0	0.1
	CI+	0.2	0.3	0.2	0.5	0.9	0.3	0.2	0.2	0.4
	CI-	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0
Poppers ³	%	0.4	0.5	0.3	0.9	1.2	0.5	0.1	0.0	0.0
	CI+	0.6	0.9	0.5	1.3	2.1	1.0	0.3	0.2	0.3
	CI-	0.3	0.3	0.2	0.6	0.7	0.2	0.0	0.0	0.0
Sedatives, Tranquillisers, Anti-depressants	%	7.7	5.5	9.9	5.1	3.9	6.3	7.3	10.5	13.4
	CI+	9.3	7.0	12.0	6.6	5.4	8.9	9.6	13.4	16.0
	CI-	8.0	5.3	10.2	4.9	3.0	6.4	7.2	10.0	12.3

¹ For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

² Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, naps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

³ Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.

Prevalence Tables – Island of Ireland

Table 3.3: Island of Ireland – Last Month Prevalence (%)

	All Adults			Young Adults						
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Any illegal drugs ¹	%	3.1	4.6	1.6	5.5	7.5	3.4	1.8	0.6	0.3
	CI+	3.6	5.7	1.9	6.7	9.8	4.4	2.5	1.2	0.8
	CI-	2.6	3.9	1.1	4.6	5.9	2.5	1.3	0.2	0.1
Cannabis	%	2.7	3.9	1.4	4.7	6.3	3.0	1.8	0.6	0.2
	CI+	3.2	5.0	1.7	5.8	8.5	4.0	2.5	1.2	0.6
	CI-	2.3	3.4	0.9	3.9	4.8	2.2	1.3	0.2	0.0
Heroin	%	0.0	0.0	0.0	0.1	0.2	0.0	0.0	0.0	0.0
	CI+	0.1	0.2	0.2	0.3	0.6	0.2	0.2	0.2	0.3
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Methadone	%	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.0	0.0
	CI+	0.2	0.2	0.3	0.3	0.5	0.5	0.4	0.4	0.3
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other opiates ²	%	1.3	1.0	1.6	1.0	0.7	1.4	1.7	1.5	1.2
	CI+	2.1	1.8	2.7	1.9	1.5	2.8	3.1	2.8	2.7
	CI-	1.5	1.0	1.8	1.1	0.4	1.5	1.7	1.4	1.2
Cocaine (total including crack)	%	0.3	0.5	0.0	0.5	0.7	0.4	0.1	0.0	0.0
	CI+	0.4	0.8	0.1	0.9	1.4	0.8	0.3	0.2	0.3
	CI-	0.1	0.2	0.0	0.2	0.2	0.1	0.0	0.0	0.0
Crack	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	CI+	0.0	0.1	0.1	0.1	0.3	0.2	0.2	0.2	0.3
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cocaine powder	%	0.3	0.5	0.0	0.5	0.7	0.4	0.1	0.0	0.0
	CI+	0.4	0.8	0.1	0.9	1.4	0.8	0.3	0.2	0.3
	CI-	0.1	0.2	0.0	0.2	0.2	0.1	0.0	0.0	0.0
Amphetamines	%	0.2	0.3	0.1	0.3	0.5	0.1	0.1	0.0	0.1
	CI+	0.3	0.6	0.2	0.6	1.2	0.4	0.3	0.4	0.5
	CI-	0.1	0.1	0.1	0.1	0.2	0.0	0.0	0.0	0.0

	All Adults			Young Adults						
	15-64	Males	Females	15-34	15-24	25-34	35-44	45-54	55-64	
Ecstasy	%	0.4	0.7	0.1	0.8	1.2	0.4	0.0	0.0	0.0
	CI+	0.6	1.1	0.2	1.3	2.1	0.8	0.3	0.2	0.3
	CI-	0.3	0.4	0.0	0.5	0.7	0.1	0.0	0.0	0.0
LSD	%	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
	CI+	0.1	0.2	0.1	0.2	0.3	0.4	0.2	0.2	0.3
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Magic mushrooms	%	0.0	0.1	0.0	0.0	0.1	0.0	0.0	0.0	0.1
	CI+	0.1	0.3	0.1	0.2	0.5	0.2	0.3	0.2	0.4
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Solvents	%	0.0	0.1	0.0	0.1	0.1	0.0	0.0	0.0	0.1
	CI+	0.1	0.2	0.1	0.2	0.5	0.2	0.2	0.2	0.4
	CI-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Poppers ³	%	0.1	0.1	0.1	0.2	0.4	0.0	0.0	0.0	0.0
	CI+	0.2	0.4	0.2	0.5	1.0	0.3	0.2	0.2	0.3
	CI-	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0
Sedatives, Tranquillisers, Anti-depressants	%	5.7	4.2	7.1	3.1	2.2	4.0	5.0	8.9	11.0
	CI+	7.0	5.5	8.9	4.2	3.3	5.9	6.9	11.7	13.6
	CI-	5.8	4.0	7.3	2.8	1.4	3.9	4.9	8.5	10.2

¹ For this study, 'any illegal drugs' refers to amphetamines, cannabis, cocaine powder, crack, ecstasy, heroin, LSD, magic mushrooms, poppers and solvents.

² Other opiates, i.e. opium, Temgesic®, buprenorphine®, Diconal®, naps, MSTs®, pethidine, DF118® (dihydrocodeine) and morphine.

³ Poppers, i.e. amyl or butyl nitrite.

All prevalence rates are based on weighted data. All figures are rounded to the nearest decimal place.



DAIRU
Drug and Alcohol
Information
and Research Unit

Department of
Health, Social Services
and Public Safety
www.dhsspsni.gov.uk
AN RIALT
Sláinte, Seirbhísí Sóisialta
agus Sébháilteachta Poiblí
MINISTERSTEW
Poblacht, Resydénter Heisín
an Fowk Siecar


Ipsos MORI

 *Northern Ireland*
**Statistics &
Research** agency


NDP
National Development Plan 2007 - 2013
Transforming Ireland