

 **Drug and Alcohol FINDINGS** Your selected document

This entry is our account of a study selected by Drug and Alcohol Findings as particularly relevant to improving outcomes from drug or alcohol interventions in the UK. Unless indicated otherwise, permission is given to distribute this entry or incorporate passages in other documents as long as the source is acknowledged including the web address <http://findings.org.uk>. The original study was not published by Findings; click on the [Title](#) to obtain copies. Free reprints may also be available from the authors – click [prepared e-mail](#) to adapt the pre-prepared e-mail message or compose your own message. Links to source documents are in [blue](#). Hover mouse over [orange](#) text for explanatory notes. The Summary is intended to convey the findings and views expressed in the study. Below are some comments from Drug and Alcohol Findings.

Open [home page](#). Get free [e-mail alerts](#) about new studies. Search studies by [topic](#) or [free text](#)

---

► [Reducing the harm from adolescent alcohol consumption: results from an adapted version of SHAHRP in Northern Ireland.](#)



DOWNLOAD PDF  
for saving to  
your computer

McKay M.T., McBride N.T., Sumnall H.R. et al.

*Journal of Substance Use*: 2012, 17(2), p. 98–121.

Unable to obtain a copy by clicking title? Try asking the author for a reprint by adapting this [prepared e-mail](#) or by writing to Dr McKay at [Michael.McKay@liverpool.ac.uk](mailto:Michael.McKay@liverpool.ac.uk). You could also try this [alternative](#) source.

*As in Australia, an alcohol harm reduction curriculum adapted for secondary schools in Northern Ireland curbed the growth in alcohol-related problems and also meant pupils drank less. Results suggest this approach might offer a more fruitful focus for education about commonly used substances than simply promoting non-use.*

**Summary** Alcohol harm reduction approaches aim to decrease the harmful consequences of drinking without requiring abstinence. School-based substance use education programmes in the United Kingdom have mainly tried to delay the onset of use, though more recent programmes have included harm reduction components. Advantages of harm reduction approaches for adolescent pupils may include not stigmatising younger drinkers, not presenting drinking as a moral issue, and being able to tailor education to the specific risk factors of the particular pupil population. Such approaches seem most relevant at the ages when young people are first drinking unsupervised by adults and experiencing intoxication.

Developed and first evaluated in Australia, the School Health and Alcohol Harm Reduction Project (SHAHRP) is an example of a harm reduction education intervention. It combines harm reduction principles with skills training, education and activities designed to encourage positive behavioural change to curb the growth in [harms](#) experienced as a result of drinking. Just such an effect was found in the [original evaluation](#), in which the number of harms [experienced](#) by pupils in SHAHRP schools was substantially and significantly less than among pupils in schools not running the lessons, and remained so at the last follow-up 17 months after lessons had ended.

Given the prevalence of underage drinking in Northern Ireland and the associated problems, it was decided to adapt SHAHRP in a culturally appropriate and curriculum-consistent manner for Northern Irish secondary (or 'high') schools. As in the original study, the adapted version was delivered over two school years in two phases. The six lessons of phase one took place when pupils were in year 10 (age 13–14), and the four in phase two the following school year. A pilot study had found that both pupils and teachers felt the programme was easy to deliver, project materials helpful and easy to follow, and activities and discussions relevant and appropriate.

Having established its feasibility, a new study starting in 2005 recruited 29 secondary schools in the Greater Belfast to test the programme's effectiveness. Nine were allocated to carry on with the normal Northern Irish alcohol education curriculum (the **control** schools), the remainder to also implement SHAHRP. In eight of these schools SHAHRP was delivered by the schools' own teachers after being trained, in 12 by local voluntary-sector drug and alcohol educators. Rather than being assigned at random, schools were assigned to the three alcohol education options so that they would be comparable in terms of gender, socio-economic profile and location. In all 2349 pupils were surveyed at the start of the study, again the following two years after the first and second phases of SHAHRP, and finally in March 2008 when lessons had ended at least 11 months before, at which time 2048 of the 2349 could be re-surveyed. Though surveys were anonymous and confidential, identifiers could be used to track changes in each individual pupil across the three years. By the end of the study pupils averaged 16½ year of age. About 60% were girls. At the start of the study 17% had not drunk alcohol and around half had drunk without adult supervision.

## Main findings

Generally the trends in how pupils drank and the harms they experienced were most favourable when SHAHRP lessons had been delivered by external specialists, next most favourable when they had delivered by the schools' own teachers, and least favourable when SHAHRP had not been implemented at all. Selected more detailed findings below.

Each survey included questions about the resultant harms pupils who had drunk at some time during the study had experienced over the past year, such as drinking more than they had planned, being sick after drinking, having hangover symptoms, being unable to remember what had happened while drunk, becoming verbally and/or physically abusive, and trouble with parents or police. Pupils divided in to four characteristic trajectories over the years of the study. Compared to those in control schools, pupils offered the SHAHRP lessons were more likely to have experienced virtually no harms during the study or a relatively low and stable level, rather than increasing and high levels of harm. When SHAHRP lessons had been delivered by external specialists, pupils were more likely to have experienced virtually no harms than when delivered by the schools' own teachers. However, both types of SHAHRP delivery significantly improved on usual lessons only.

The drinkers among the pupils were also asked how much they had drunk last time. On this measure pupils again divided in to four characteristic trajectories. Compared to those in control schools, at each follow-up pupils offered the SHAHRP lessons were more likely say they had drunk very little than to have reported increasing and by the end of the study relatively high levels of drinking. When the lessons had been delivered by external specialists, pupils were more likely to consistently have drunk relatively little than when

delivered by the schools' own teachers.

Each survey also included questions about the harms pupils had experienced over the past year arising from someone else's drinking, such as verbal or physical abuse, sexual harassment, or damage to personal property. Compared to those in control schools, pupils offered the SHAHRP lessons were least likely to have experienced a steep rise in such harms ending in relatively high levels. Whether SHAHRP lessons had been delivered by external specialists or the schools' own teachers did not significantly affect the trends.

Pupils offered SHAHRP lessons were more likely than those in control schools to have become more knowledgeable about alcohol over the study and to end with relatively high levels of knowledge, most so when the lessons had been delivered by external specialists. However, both types of SHAHRP delivery significantly improved on usual lessons only. Results were similar in respect of developing safer attitudes to drinking.

### The authors' conclusions

A [research review](#) associated with [guidance](#) on alcohol education from the National Institute for Health and Clinical Excellence remarked that the Australian SHAHRP evaluation offered evidence that programmes focusing on harm reduction through skills-based activities can produce medium to long term reductions in alcohol use and in particular, risky drinking behaviours. However, the review queried the transferability of these programmes and results to the UK. The featured study shows that in the UK too, classroom-based harm reduction education can have a significant impact on the harm adolescents experienced from drinking. The research also suggests these lessons need to incorporate interactive learning, start just prior to and during the times when pupils first try drinking, be culturally sensitive, and provide realistic scenarios and deal with realistic issues.

Compared to control schools, pupils in SHAHRP schools were significantly more likely to be among groups characterised by better growth in knowledge about alcohol and its effects, safer alcohol-related attitudes, fewer harms from one's own and other's drinking, and less alcohol consumption. These differences were maintained over an 11-month period after the lessons had ended, though in some cases with diminished strength. External facilitation of the lessons was associated with the most desirable outcomes, particularly with respect to knowledge and attitudes, harms from one's own drinking, and alcohol consumption.

SHAHRP offers abstainers, novice drinkers and more experienced drinkers alike the opportunity to reflect on use, harm and personal safety, including the importance of trusted friends, basic first aid techniques, group transport home, mobile phone availability, not to make decisions while drunk, identify friends becoming drunk, drink-spiking, mixing substances, and arguments and aggressive behaviour. The results show that young people are capable of processing harm reduction messages developed and presented within the reality of their drinking experiences. SHAHRP addresses harms without causing any increase in drinking (in fact, the reverse) or decreasing rates of abstinence.

It was unfortunate that two of the schools allocated to the control group withdrew from the study, partially upsetting the attempt to ensure comparability of the schools operating the three alcohol education options. However, differences were adjusted for statistically. Also, no systematic record was kept on the alcohol

education delivered to control subjects. In Northern Ireland this typically is embedded in the curriculum as part of science or citizenship lessons, so would be identical to that received by intervention students.

**FINDINGS** Together with the original Australian evaluation, this UK study represents fairly strong evidence that if it focuses on this task, a school curriculum can reduce drink-related problems. In [Australia](#) harm-reduction effects were greatest among the higher risk pupils who had already drunk without adult supervision; at each follow-up point they experienced about 20% fewer harms than control pupils.

In that study too, though still very much in the minority, by the last follow up there were a third more abstainers among SHAHRP than control pupils. By the end of the featured study about 6% of control pupils had never drunk alcohol compared to 6% of SHAHRP pupils taught by external staff and 3.5% taught by their teachers. These findings offer little support to concerns that safer drinking lessons will encourage more pupils to drink.

In the more restrictive youth drinking environment of the USA, a programme forefronting alcohol problem reduction among its aims has [produced similar findings](#) to that in Australia. It retarded growth in alcohol problems (such as getting drunk or sick or complaints from parents and friends), but only among pupils who had already drunk without adult supervision, and only if the lessons did not occur too early to coincide with the development of this drinking pattern. After disappointing initial results, another US substance use education programme including alcohol adopted harm reduction objectives. The revised programme [resulted](#) in a significant reduction in risky or harmful drinking. Parallel and consistent findings in different countries with different curricula suggests that harm reduction education on drinking has a real and transferable impact in Western drinking cultures. Such findings contrast with unconvincing evidence from trials of [substance use education](#) in general and [alcohol education](#) in particular.

For the UK the most important [guidance](#) on alcohol education was issued in 2007 by the National Institute for Health and Clinical Excellence. It said this "should aim to encourage children not to drink, delay the age at which young people start drinking and reduce the harm it can cause among those who do drink". Recommendations included ensuring alcohol education is an integral part of the science and PSHE curricula.

[Inspections](#) in 2012 of PSHE lessons suggest English schools are far from adequately implementing this recommendation, in particular in respect of education aimed at reducing alcohol-related harm. In just under half the inspected schools had pupils learnt how to keep themselves safe in a variety of situations, and the deficits were particularly noticeable in respect of drinking. Inspectors found that although pupils understood the dangers to health of tobacco and illegal drugs, they were far less aware of the physical and social damage associated with risky drinking. Some did not know the strength of different alcoholic drinks or make the links between excessive drinking and issues such as heart and liver disease and personal safety. The report attributed these deficiencies in part to inadequacies in subject-specific training and support for PSHE teachers, particularly in teaching sensitive and controversial topics.

Last revised 13 June 2013. First uploaded 12 June 2013

- ▶ [Comment on this entry](#)
- ▶ [Give us your feedback on the site \(one-minute survey\)](#)
- ▶ Open [home page](#) and enter [e-mail address](#) to be alerted to new studies

**Top 10 most closely related documents on this site. For more try a [subject or free text search](#)**

[School lessons reduce alcohol-related harm](#) STUDY 2004

[Effects of a school-based prevention program on European adolescents' patterns of alcohol use](#) STUDY 2011

[Initial outcomes from Australian alcohol harm reduction curriculum for secondary schools](#) STUDY 2001

[Does successful school-based prevention of bullying influence substance use among 13- to 16-year-olds?](#) STUDY 2010

[The effectiveness of a school-based substance abuse prevention program: 18-month follow-up of the EU-Dap cluster randomized controlled trial](#) STUDY 2010

[Substance use outcomes 5½ years past baseline for partnership-based, family-school preventive interventions](#) STUDY 2008

[Everyone is NOT doing it - important prevention message for early teens](#) STUDY 2000

[Substances, adolescence \(meta-analysis\)](#) STUDY 2003

[Project SUCCESS' effects on the substance use of alternative high school students](#) STUDY 2010

[School programme successfully revised to focus more on harm reduction](#) STUDY 2005