

Factsheet: Sedatives and tranquillisers - the Irish situation

August 2023




Table of Contents

Table of Contents 1

Glossary of terms..... 1

What are sedatives and tranquillisers? 2

What do sedatives and tranquillisers do? 2

How do we know how many people use sedatives or tranquillisers in Ireland? 2

How many people use sedatives or tranquillisers in Ireland? 2

Use among young people..... 3

How many people receive treatment for sedative and tranquilliser use? 3

How many people die from misuse of sedatives and tranquillisers? 4

Non-fatal overdoses and drug-related emergencies 4

What does the law say about sedatives and tranquillisers? 4

Seizures of sedatives and tranquillisers 5

References..... 5

Glossary of terms

Last month prevalence – refers to the proportion of the sample that reported using a named drug in the 30-day period prior to the survey. Last month prevalence is often referred to as current use. A proportion of those reporting current use may be occasional (or first-time) users who happen to have used in the period leading up to the survey. It should therefore be noted that current use is not synonymous with regular use.

Last year prevalence – refers to the proportion of the sample that reported using a named drug in the year prior to the survey. Last year prevalence is often referred to as recent use.

Lifetime prevalence – refers to the proportion of the sample that reported ever having used the named drug at the time they were surveyed. A person who records lifetime prevalence may or may not be currently using the drug. Lifetime prevalence should not be interpreted as meaning that people have necessarily used a drug over a long period of time or that they will use the drug again in future.

Prevalence – refers to the proportion of a population that has used a drug over a particular time period.

Statistically significant – a result is deemed statistically significant if it is unlikely to have occurred by chance, and hence provides enough evidence to reject the hypothesis of ‘no effect’. As used in statistics, ‘significant’ does not mean important or meaningful. A small, but important, real-world difference may fail to reach significance in a statistical test, while a statistically significant finding may have no practical consequence.

What are sedatives and tranquillisers?

'Sedatives' and 'tranquillisers' are commonly used terms for a group of medicines which depress, slow down or calm the brain and central nervous system. ¹

What do sedatives and tranquillisers do?

Sedatives and tranquillisers are commonly used terms for the same group of medicines which depress, slow down, or calm the brain and central nervous system. They are mainly benzodiazepines ('benzos'), but other drugs with the same effects (e.g. zolpidem and zopiclone ('Z-drugs')) are included in this group. Medically they are often referred to as hypnotics, which induce sleep, and as anxiolytics or anti-anxiety agents. ¹

How do we know how many people use sedatives or tranquillisers in Ireland?

Every four years a survey of the general population takes place to estimate the number of people in Ireland who use alcohol and other drugs. ¹ Face-to-face interviews take place with respondents aged 15+ normally resident in households in Ireland. This type of survey is not designed to include people who do not normally live in private households, such as prisoners or hostel dwellers. ³ (For other populations, see our [prevalence sources](#).)

Drug prevalence surveys were undertaken in 2002–03, 2006–07, 2010–11, and 2014–15. These surveys were commissioned on an all-island basis by the National Advisory Committee on Drugs and Alcohol in the Republic of Ireland, and the Department of Health, Social Services and Public Safety in Northern Ireland. The most recent survey 2019–20 National Drug and Alcohol Survey (NDAS), which took place in the Republic of Ireland, was managed by the Health Research Board.

How many people use sedatives or tranquillisers in Ireland?

The 2019–20 NDAS survey involved 5,762 people in Ireland. ¹ The results for Ireland showed that:

- 12.7% of respondents had ever used sedatives/tranquillisers in their lifetime, corresponding to 499,000 of the general population in Ireland aged 15 years and older.
- 5.5% of respondents (213,000 of the general population) and 3.2% of respondents (125,000 of the general population) had used sedatives/tranquillisers in the last year and last month, respectively.
- 0.5% of respondents (19,000 of the general population) had used sedatives/tranquillisers in a non-medical way in the last 12 months.
- There was a statistically significant decrease in recent use reported by all adults (from 7.1% in 2014–15 to 5.5% in 2019–20).
- The average age of first sedative/tranquilliser use was 36.7 years (median: 34 years).
- The average age of respondents who reported recent sedative/tranquilliser use was 50.9 years (median: 49 years).

Females were more likely than males to report recent use of sedatives/tranquillisers (6.1% versus 4.8%). Those aged 65 years and older were more likely than younger age groups to report recent sedative/tranquilliser use (8.0%). Females aged 65 years and older had the highest prevalence of recent use (11.5%). Of those who had used sedatives/ tranquillisers in the last year, 87.2% stated that all were prescribed, 10.2% stated that none were prescribed, and 2.7% stated that some were prescribed while others were not.¹

Use among young people

The [European School Survey Project on Alcohol and Other Drugs](#) (ESPAD) collects comparable data on substance use among 15–16-year-old students in 35 countries. According to the 2020 ESPAD report, 2.6% of respondents had taken tranquillisers without a prescription.²

For further information on sedative and tranquilliser use (prevalence and treatment) by County see our [Regional data factsheets](#).

How many people receive treatment for sedative and tranquilliser use?

The [National Drug Treatment Reporting System](#) (NDTRS) provides data on treated drug and alcohol misuse in Ireland. The most recent published data from the NDTRS³ shows that:

Z-drugs (non-benzodiazepine hypnotic sedative drugs such as zolpidem, zopiclone) as a main problem continued to account for 0.8% of cases in 2022, a decrease from 1.1% in 2016.

Benzodiazepines were the main problem for 10.7% of cases in 2022, an increase from 9.7% in 2016. 374 (8.4%) of these were new cases. Benzodiazepines (32.2%) were the fourth most common additional drug reported among cases with polydrug use in 2022.

Table 2: Main problem drug (excluding alcohol) reported in 30 days prior to treatment, NDTRS 2016 to 2022

| | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 |
|---------------------------------|------------|------------|------------|-------------|-------------|-------------|-------------|
| | N (%) | N (%) | N (%) | N (%) | N (%) | N (%) | N (%) |
| All cases | 9227 | 8922 | 10274 | 10664 | 9702 | 10769 | 12009 |
| Benzodiazepines | 897 (9.7) | 868 (9.7) | 999 (9.7) | 1082 (10.1) | 1097 (11.3) | 1218 (11.3) | 1287 (10.7) |
| Z-drugs | 103 (1.1) | 82 (0.9) | 48 (0.5) | 72 (0.7) | 72 (0.7) | 80 (0.7) | 98 (0.8) |
| New cases | 3626 | 3257 | 3962 | 3979 | 3796 | 4206 | 4456 |
| Benzodiazepines | 353 (10.0) | 290 (8.9) | 345 (8.7) | 340 (8.5) | 392 (10.3) | 418 (9.9) | 374 (8.4) |
| Z-drugs | 41 (1.2) | 22 (0.7) | 17 (0.4) | 24 (0.6) | 27 (0.7) | 26 (0.6) | 15 (0.3) |
| Previously treated cases | 5335 | 5242 | 5872 | 5927 | 5441 | 6090 | 6860 |
| Benzodiazepines | 500 (9.4) | 534 (10.2) | 596 (10.1) | 661 (11.2) | 631 (11.6) | 742 (12.2) | 829 (12.1) |
| Z-drugs | 58 (1.1) | 56 (1.1) | 31 (0.5) | 43 (0.7) | 42 (0.8) | 54 (0.9) | 79 (1.2) |

See the NDTRS report³ for more information on polydrug use - additional problem drugs for all cases and new cases.

For more detailed treatment data (age, gender, employment status) up to 2021 see library [interactive tables](#).

How many people die from misuse of sedatives and tranquillisers?

The [National Drug-Related Deaths Index](#) (NDRDI) is a database of cases of death by drug and alcohol poisoning, and deaths among drug users and people who are alcohol dependent. In 2020, Almost 6 in 10 (228) poisoning deaths involved benzodiazepines and many of them had more than one type of benzodiazepine (including diazepam, flurazepam, alprazolam). Diazepam poisoning deaths have decreased slightly from 133 in 2011 to 130 in 2020. Almost 6 in 10 poisoning deaths involved other prescription drugs, most commonly antidepressants and antiepileptics⁴

Table 1: Benzodiazepines and other prescription drugs implicated in poisoning deaths, NDRDI 2011-2020

| | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|------|------|------|------|------|------|------|------|------|------|
| All poisoning deaths | 314 | 281 | 341 | 319 | 327 | 314 | 340 | 354 | 371 | 409 |
| Drug group: | | | | | | | | | | |
| Any benzodiazepine implicated | 170 | 129 | 163 | 164 | 147 | 147 | 150 | 196 | 186 | 228 |
| Any other prescription drug implicated** | 137 | 138 | 184 | 193 | 189 | 202 | 179 | 191 | 199 | 227 |
| Individual drug: | | | | | | | | | | |
| Diazepam | 133 | 92 | 113 | 121 | 109 | 100 | 97 | 122 | 102 | 130 |
| Flurazepam | 50 | 30 | 42 | 36 | 35 | 43 | 35 | 47 | 30 | 30 |
| Poly drug poisonings: | | | | | | | | | | |
| More than one benzodiazepine implicated | 70 | 41 | 62 | 61 | 62 | 71 | 72 | 96 | 68 | 89 |
| More than one other prescription drug implicated** | 48 | 53 | 76 | 96 | 94 | 102 | 98 | 93 | 97 | 119 |

** Most commonly antidepressants and antiepileptics. Also includes non-benzodiazepine sedatives

Non-fatal overdoses and drug-related emergencies

According to the [Hospital In-Patient Enquiry Scheme](#) (HIPE), there were 5,402 cases of non-fatal overdose discharged from Irish hospitals in 2020. There was evidence of benzodiazepines in 1,111 of cases.⁵

What does the law say about sedatives and tranquillisers?

Under the Medicinal Products (Prescription and Control of Supply) Regulations 2003–2008, a prescription medication can only be supplied in accordance with a prescription, and the supply must be made from a registered pharmacy by or under the personal supervision of a registered pharmacist. It is illegal for prescription medicines to be supplied through mail-order or internet sites. A person who has in his possession a prescription medicine containing a substance controlled under

the misuse of drugs legislation for the purpose of selling or otherwise supplying it is guilty of an offence under that legislation.

Changes to regulations under the [Misuse of Drugs Amendment Act 2016](#) introduced stricter controls on benzodiazepines and an initiative to tackle overprescribing. You can find more information about Irish drug laws, offences and penalties on the [Citizens Information Board website](#).

Seizures of sedatives and tranquillisers

Following a peak in 2013 (N=861), the number of seizures of hypnotic and sedative substances decreased annually until 2016. A 73% increase in these seizures was recorded between 2016 and 2017, followed by a 49% decrease between 2017 and 2018. However, the number of seizures analysed in 2019 (N=1269) was more than four times higher than those reported in 2018 (N=309). Increases were evident between 2019 and 2020 (8%) and again between 2020 and 2021 (6%).⁶

References

1. Mongan D, Millar SR, and Galvin B (2021) [The 2019–20 Irish National Drug and Alcohol Survey: main findings](#). Dublin: Health Research Board.
2. Sunday S, Keogan S, Hanafin J and Clancy L (2020) [ESPAD 2019 Ireland](#). Dublin: TobaccoFree Research Institute Ireland.
3. O'Neill D, Lyons S and Carew AM (2023) [National drug treatment reporting system, 2022 drug treatment demand](#). Dublin: Health Research Board.
4. Health Research Board (2023) [Health Research Board reports latest drug-related deaths figures](#). Dublin: Health Research Board.
5. Millar, S (2022) [Not-fatal drug-related hospital admissions in Ireland, 2020](#). Drugnet Ireland, Issue 83, Winter 2022, pp. 16-20
6. Health Research Board. Irish National Focal Point to the European Monitoring Centre for Drugs and Drug Addiction (2023) [Ireland: national report for 2022 - drug markets and crime](#). Dublin: Health Research Board.

Further reading:

European Monitoring Centre for Drugs and Drug Addiction. (2018) [Perspectives on drugs: the misuse of benzodiazepines among high-risk opioid users in Europe](#). Lisbon: EMCDDA

For regional Irish data, please see our [regional factsheets webpage](#).

How to cite this factsheet:

HRB National Drugs Library (2023) Sedatives and tranquillisers - the Irish situation. Dublin: HRB National Drugs Library www.drugsandalcohol.ie/24954

Other factsheets in this series:

- [Alcohol - the Irish situation](#)
- [Cannabis - the Irish situation](#)
- [Cocaine - the Irish situation](#)
- [MDMA/ecstasy - the Irish situation](#)
- [New psychoactive substances \(NPS\) - the Irish situation](#)
- [Opioids - the Irish situation](#)
- [Smoking - the Irish situation](#)
- [Young people, drugs and alcohol - the Irish situation](#)

HRB National Drugs Library – Find the evidence

www.drugsandalcohol.ie

- Freely available collection of Irish and international drug and alcohol research
- Quick updates – e-newsletter, Drugnet Ireland and Twitter (@HRBdrugslibrary)
- Key Irish data resources – treatment, deaths, prevalence, regional data, alcohol links
- Key Irish data summaries – factsheets & annual national reports
- Policy – policy webpage & Dail debates
- Practitioner portal – specific webpages for those working in the area
- Course directory
- Glossary – with explanations of terms and acronyms
- Research resources for those conducting literature searches and other research
- Search our collection – basic and advanced (you can save your results)

HRB National Drugs Library

Health Research Board
Grattan House
67-72 Lower Mount Street
Dublin 2, Ireland

e drugslibrary@hrb.ie

w www.drugsandalcohol.ie

 [@HRBdrugslibrary](https://twitter.com/HRBdrugslibrary)