

# ARAN - Access to Research at NUI Galway

Provided by the author(s) and NUI Galway in accordance with publisher policies. Please cite the published version when available.

Title	Policy priorities for girls' health	
Author(s)	Nic Gabhainn, Saoirse; Kelly, Colette; Gavin, Aoife	
Publication Date	2021-04	
Publication Information	Nic Gabhainn, Saoirse, Kelly, Colette, & Gavin, Aoife. (2021). Policy priorities for girls' health. Galway: Health Promotion Research Centre, National University of Ireland Galway. DOI: 10.13025/xsew-yf04	
Publisher	Health Promotion Research Centre, National University of Ireland Galway	
Link to publisher's version	https://doi.org/10.13025/xsew-yf04	
Item record http://hdl.handle.net/10379/16810		
DOI	http://dx.doi.org/10.13025/xsew-yf04	

Downloaded 2022-08-05T09:09:13Z

Some rights reserved. For more information, please see the item record link above.





## **Key Policy Priorities for Girls Health**

Saoirse Nic Gabhainn, Colette Kelly and Aoife Gavin, HBSC Ireland Research Team Health Promotion Research Centre, National University of Ireland Galway

April 2021

## Rationale for including girls

- Girls deserve health and wellbeing in their own right, from a human rights perspective, as protected by international law
- Many health issues affecting women have their roots in girlhood, effective prevention and health promotion must commence during this developmental period
- The social and economic benefits of investing in girl's health is clear and evidence based.

Childhood and adolescence are critical developmental periods with unique challenges and opportunities to influence health and wellbeing (Inchley et al., 2020a). Investment in these life stages and support for these key formative years can provide better outcomes for children, with an ecological life-course approach recommended (Tomlinson et al., 2021). Investment is particularly important to address inequalities in health. The right to health and development for children and adolescents is central to the United Nations Convention on the Rights of the Child (United Nations, 1989).

Previous European Child and Adolescent Health Strategies (WHO, 2005 and 2014) and the forthcoming strategy¹ (up to 2030) builds on these commitments with the overarching goal of promoting healthier populations. Health outcomes for children and adolescents are good in Europe, including in Ireland, but there are discrepancies in health within countries, including between boys and girls, and between girls, especially for those with social and economic disadvantage. The cycle of disadvantage can be broken by addressing inequities during childhood and adolescence to enable children and young people to reach their full potential.

#### **Key issues from policy**

In this section we draw on the 2030 Agenda for sustainable development and specifically SDG3, SDG5 and SDG10 on health and wellbeing, gender equality and reducing health inequalities, the WHO Global Strategy for Women's, Children's and Adolescents' Health 2016–2030 (WHO, 2016) the Irish National Strategy for Women and Girls 2017-2020 (Department of Justice and Equality, 2017), and the forthcoming European Strategy for Child and Adolescent Health.

<sup>&</sup>lt;sup>1</sup> We have worked on this strategy with WHO Europe, including the provision of pan-European data, and leading the direct consultation with young people on their contributions to the strategy development process.





#### **Principles and Priorities**

Human-rights based approach, equity-driven, gender-responsive, life course approach, intersectoral action, and ensuring effective participation.

- Strengthening governance for women's health and well-being, with women and girls at the centre; Embed gender equality in decision-making
- Eliminating discriminatory values, norms and practices that affect the health and well-being of girls and women; Ensure visibility, equal and active citizenship, and leadership
- Tackling the impact of gender and social, economic, cultural and environmental determinants on the health and well-being of girls and women; Advance socioeconomic equality
- Improving health system responses to the health and well-being of women and girls; Ensure cross-sectoral evidence based responses to documented priorities.

## Key health issues for girls in Ireland

In this section we focus on differences between boys and girls, and further inequalities between girls based on their personal, family or social characteristics.

The findings reported here are drawn from our work on the fifty-country WHO collaborative study, Health Behaviour in School-aged Children (HBSC): primarily the 2020 Irish National Report (Költő et al., 2020); the 2020 International HBSC report (Inchley et al., 2020) and the Irish Trends report 1998-2018 (Gavin et al., 2021). All differences reported are statistically significant. Further information and all outputs from the HBSC Ireland study are available from <a href="www.nuigalway.ie/hbsc">www.nuigalway.ie/hbsc</a>. Information on the international HBSC study can be found at <a href="www.hbsc.org">www.hbsc.org</a>.

The six summaries below reflect published findings from the HBSC study, presented by health outcomes for girls both positive and negative, health behaviours for girls both positive and negative and the social determinants of health for girls. More detailed tables are presented from page four. The Irish HBSC team operate a knowledge translation helpdesk to provide evidence to policy and decision-makers. Greater detail on the findings reported, methodologies employed, and analytic approaches, or further analyses of the existing datasets, can be requested by contacting us at <a href="https://doi.org/10.1007/nbsc/">https://doi.org/10.1007/nbsc/</a>.

#### Summary of Positive Health Behaviours

Girls are less likely than boys to engage in physical activity, and there has only been a marginal improvement in this over time. Girls are less likely than boys to eat breakfast. Physical activity and breakfast eating are least common among girls with social or economic disadvantage. See table 1 below for more detail.

#### Summary of Negative Health Behaviours

Girls are more likely than boys to report being cyberbullied, being on a diet and eating sweets at least daily. Dieting, bullying and fighting are more frequent among girls with





social or economic disadvantage. Condom use among girls has decreased over time, while dieting has increased. See table 2 below for more detail.

## Summary of Positive Health Outcomes

Girls are less likely than boys to report high life satisfaction, self-reported happiness, self-rated health and good mental health. There is evidence of a reduction over time in the wellbeing of girls. Girls living with social economic disadvantage have lower wellbeing. See table 3 below for more detail.

## Summary of Negative Health Outcomes

Girls are more likely than boys to report frequent somatic and psychological symptoms. There is evidence of increases over time in symptoms. Girls living with social or economic disadvantage report more frequent symptoms. Girls are more likely than boys to report that they are too fat, and this is even more common in girls from low affluent families. Some sub-groups of girls also report elevated rates of medically attended injuries. See table 4 below for more detail.

## Summary of the Social Determinants of Health

Girls are more likely than boys to report intense and problematic social media use, high levels of pressure from schoolwork, and feeling discriminated against on the basis of being a girl and due to their age. Girls are less likely than boys to find it easy to talk to their fathers, to feel safe in the area that they live or feel they can ask neighbours for help. Schoolwork pressure has increased over time, while feeling safe and asking neighbours for help has reduced. Girls living with social or economic disadvantage report fewer close friends, less ease of communication with friends and parents, more intense social media use, and are less likely to like school or feel supported or accepted by their classmates. See table 5 below for more detail.





# **Table 1. Positive Health Outcomes**

Issue	Pattern	Some girls at more risk than others	Source
Life	Girls have lower life satisfaction than boys;	Girls from low affluence families and lower social class	Költő et al. (2020)
satisfaction	Life satisfaction in girls has decreased between 2002 and 2018	groups have lower life satisfaction than those from higher affluence and higher social class families; Girls from the Traveller community report lower life satisfaction than their peers	Molcho et al. (2008)
Self-rated health	Girls have lower self-rated health than boys	Girls from low affluence and lower social class families have lower self-rated health than those from higher affluence and higher social class families	Költő et al. (2020) Inchley et al. (2020b)
Mental health	Girls have poorer mental wellbeing than boys as assessed by the Mental Health Inventory and by the WHO-5	Girls from lower social class families have poorer mental wellbeing than those from higher social class families	Költő et al. (2020)
Self-reported happiness	Girls have lower rates of self-reported happiness than boys. Happiness in girls has decreased between 1998 and 2018	Girls from immigrant families have lower rates of self-reported health than their peers.	Költő et al. (2020) Molcho et al. (2008)

# **Table 2. Negative Health Outcomes**

Issue	Pattern	Some girls at more risk than others	Source
Symptoms	Headache, stomach-ache, sleep difficulties, feeling	Girls from the Traveller community, immigrant families, and	Inchley et al. (2020b)
	nervous, feeling low, feeling irritable and feeling	DEIS schools report more frequent headaches than their	Molcho et al. (2008)
	dizzy are all more frequent among girls than	peers;	
	among boys by age 15;	Girls from low affluence families are more likely to report	
	Frequent headaches and feeling low is higher for	2+ symptoms per week than those from higher affluence	
	girls in 2018 than in 1998	families	
Feeling fat	Girls are more likely than boys to report that they	Girls from low affluence families are more likely to report	Inchley et al. (2020b)
	are too fat at ages 11, 13 and 15	that they are too fat than those from higher affluence	
		families	
Injuries	Girls are less likely to be injured requiring medical	Girls who live with a disability or chronic illness are more	Inchley et al. (2020b)
	attention than boys	likely than other girls to have been injured requiring	Molcho et al. (2008)
		medical attention;	
		Girls from low affluence families are more likely to be	
		injured requiring medical attention than those from higher	
		affluence families	





**Table 3. Positive Health Behaviours** 

Issue	Pattern	Some girls at more risk than others	Source
Physical	Girls report less frequent moderate	Girls from low affluence and lower social class families are less likely to	Inchley et al. (2020b)
Activity	and vigorous physical activity than	report that frequently engage in physical activity than those from higher	Költő et al. (2020)
	boys; Physical activity among girls has	affluence or higher social class families	Gavin et al. (2021)
	increased only marginally between		
	1998 and 2018		
Eating	Girls are less likely to report eating	Girls from low affluence and lower social class families are less likely to	Költő et al. (2020)
Breakfast	breakfast than boys	report eating breakfast, than those from higher affluence or higher	Molcho et al. (2008)
		social class families;	Inchley et al. (2020b)
		Girls from the Traveller community, those from immigrant families and	
		those in DEIS schools are less likely to report eating breakfast than their	
		peers	

# **Table 4. Negative Health Behaviours**

Issue	Pattern	Some girls at more risk than others	Source
Cyberbullying	Girls are more likely than boys to be	Girls from low affluence and lower social class families are more likely	Inchley et al. (2020b)
	cyberbullied	to report that they are cyberbullied than those from higher affluence or	Költő et al. (2020)
		higher social class families	
Bullying		Girls from the Traveller community, and those who live with a disability	Molcho et al. (2008)
		or chronic illness report higher levels of traditional face-to-face bullying	
		than their peers	
Fighting		Girls who live with a disability or chronic illness and those attending	Molcho et al. (2008)
		DEIS schools are more likely than other girls to be involved in a physical	
		fight	
Condom use	In 2018 fewer girls report condom use		Gavin et al. (2021)
	at last intercourse than in 2010 or		
	2014.		
Dieting or	Girls are more likely to reporting being	Girls from lower social class families are less likely to report dieting than	Inchley et al. (2020b)
doing	on a diet than boys;	those from higher social families;	Költő et al. (2020)
something	Girls being on a diet has increased over	Girls from the Traveller community, from immigrant families, and those	Gavin et al. (2021)
else to lose	time between 2002 and 2018,	with a disability or chronic illness are more likely to report dieting than	Molcho et al. (2008)
weight	especially among younger girls	their peers	
Sweets	Girls are more likely than boys to eat		Költő et al. (2020)
	sweets daily or more frequently		





**Table 5. Social Determinants of Health** 

Issue	Pattern	Some girls at more risk than others	Source
Friends		Girls from the Traveller community and from immigrant	Molcho et al. (2008)
		families, and those with a disability or chronic illness are less	Inchley et al. (2020b)
		likely to have 3+ friends of the same sex than their peers;	
		Girls from low affluence families are less likely to find it easy to	
		talk to their peers about things that really bother them than	
		those from higher affluence families	
Social Media	Girls report higher intensity of social media use	Girls from low affluence and low social class families report	Inchley et al. (2020b)
	than boys at age 13 and 15;	higher intensity social media use than those from higher	Költő et al. (2020)
	Girls report higher levels of problematic social	affluence and social class families.	
	media use than boys at ages 11 and 13		
Families	Girls are less likely than boys to find it easy to	Girls from low affluence families are less likely to report that it	Inchley et al. (2020b)
	talk to their fathers about things that really	is easy to talk to their fathers and their mothers about things	
	bother them at ages 11, 13 and 15	that really bother them than those from higher affluence	
		families	
School	At age 15, girls are more likely than boys to	Girls from the Traveller community and from immigrant	Gavin et al. (2021)
	report being pressured by schoolwork;	families, and those with a disability or chronic illness are less	Molcho et al. (2008)
	There have been substantial increases in	likely to like school or feel accepted by other students than	Inchley et al. (2020b)
	schoolwork pressure reported by girls between	their peers;	
	1998 and 2018.	Girls from low affluence families are less likely to report that	
		they like school, or feel supported by classmates, than those	
		from higher affluence families.	0 1 1 (2004)
Safety	Girls are less likely to feel safe in the area that		Gavin et al. (2021)
	they live than boys;		
	Fewer girls report that they feel safe in their local area in 2002 than in 2018		
N1 - ! - l- l			Carrier at al. (2024)
Neighbours	There have been reductions in the proportion		Gavin et al. (2021)
	of girls who report they can ask neighbours for		
Diagrippin ati	help between 2002 and 2018		Malkor et al
Discrimination	Girls are more likely than boys to report being		Walker et al.
	discriminated against on the basis of being a		(forthcoming, 2021)
	girl or a boy, and on the basis of their age		





#### References

Department of Justice and Equality (2017). *Irish National Strategy for Women and Girls 2017-2020:* Creating a Better Society for All. Dublin: Department of Justice and Equality. download

Gavin, A., Költő, A., Kelly, C., Molcho, M., & Nic Gabhainn, S. (2021). *Trends in Health Behaviours, Health Outcomes and Contextual Factors between 1998-2018: findings from the Irish Health Behaviour in Schoolaged Children Study.* Dublin: Department of Health. <u>download</u>

Inchley, J., Currie, D., Budisavljevic, S., Torsheim, T., Jåstad, A., Alina Cosma, A., Kelly, C., Arnarsson, A.M., & Samdal, S. (2020b). *Spotlight on Adolescent Health and Well-being. Findings from the 2017/2018 Health Behaviour in School-aged Children (HBSC) survey in Europe and Canada. International report. Volume 2. Key data*. Copenhagen: World Health Organization Regional Office for Europe. <u>download</u>

Inchley, J.C., Stevens G.W.J.M, Samdal, O., & Currie, D.B. (2020). Enhancing understanding of adolescent health and well-Being: the Health Behaviour in School-aged Children study. *Journal of Adolescent Health, 66*, s3-s5. download

Költő, A., Gavin, A., Molcho, M., Kelly, C., Walker, L., & Nic Gabhainn, S. (2020). *The Irish Health Behaviour in School-aged Children (HBSC) Study 2018*. Dublin: Department of Health. download

Molcho, M., Kelly, C., Gavin, A., & Nic Gabhainn, S. (2008). *Inequalities in health among school-aged children in Ireland*. Dublin: Department of Health and Children. <u>download</u>

Tomlinson, M., Hunt, X., Daelmans, B., Rollins, N., Ross, D., & Oberklaid, F. (2021). Optimising child and adolescent health and development through an integrated ecological life course approach. *British Medical Journal*, 2021; 372. download

United Nations (1989) Convention on the Rights of the Child. Geneva: United Nations. download

United Nations (2015). *Transforming our world: the 2030 Agenda for Sustainable Development.* Geneva: United Nations. download

Walker, L., Gavin, A., Költő, A., Kelly, C., Molcho, M., & Nic Gabhainn, S. (2021). *HBSC Ireland 2018 full variable report: Socio-demographic pattern in health outcomes, behaviours and the social contexts of health*. Galway: Health Promotion Research Centre, NUI Galway. (with accompanying interactive data visuals, available at: <a href="http://www.nuigalway.ie/hbsc/hbscireland/2018study/">http://www.nuigalway.ie/hbsc/hbscireland/2018study/</a>)

World Health Organization (2005). *European Strategy for Child and Adolescent Health*. Copenhagen: World Health Organization. <u>download</u>

World Health Organization (2014). *Investing in Children: The European Child and Adolescent Health Strategy 2015-2020.* Copenhagen: World Health Organization. download

World Health Organization (2016). *Global Strategy for Women's, Children's and Adolescents' Health 2016–2030: Survive, Thrive, Transform.* Geneva: World Health Organization. <u>download</u>



