
Naloxone knowledge among a sample of people that regularly consume illicit stimulants in Australia, 2022

Julia Uporova, Amy Peacock and Rachel Sutherland
National Drug and Alcohol Research Centre, UNSW Sydney, Sydney, Australia



Introduction

Naloxone (Narcan, Prenoxad and Nyxoid) is an opioid antagonist used to counter the effects of overdose from opioids (1), and is available for <u>free to anyone who may experience</u>, <u>or witness</u>, <u>an opioid overdose</u>. Most research has focused on naloxone knowledge amongst people who inject drugs and/or use opioids (2), with much less known about awareness among other groups of people who use drugs who may be able to provide immediate bystander support.

Results

Using data from the 2022 <u>Ecstasy and Related Drugs Reporting System (EDRS)</u> (N=700) we found that among a sample of people who regularly use ecstasy and/or other illicit stimulants recruited from capital cities:

- Almost half (46%) reported that they had ever heard of naloxone.
- Of those that reported awareness and responded (n=312), the large majority (89%; 40% of the total sample) were able to correctly identify the purpose of naloxone (i.e., to reverse a drug overdose), whereas the remaining 11% (n=35) were unaware of its purpose or nominated an incorrect purpose (e.g., to help stop smoking).

Discussion

Awareness of naloxone among the EDRS sample was lower than has been reported among samples of people who inject drugs (84% in 2022), which is not surprising given the comparatively higher levels of opioid use among the latter (3). While people who inject drugs and/or use opioids should remain the focus of naloxone distribution programs, there may be benefit in increasing naloxone awareness and uptake among other groups of people who use drugs. This is particularly pertinent given that illicit substances such as cocaine have been found to contain opioids and have in some cases contributed to severe overdoses in Australia (4). Further, people who use illicit stimulants may be more likely to be in settings where overdoses might occur and be around people who intentionally or unintentionally use opioids.

References

- 1. National Center for Biotechnology Information (2023). PubChem Compound Summary for CID 5284596, Naloxone. Retrieved January 12, 2023 from https://pubchem.ncbi.nlm.nih.gov/compound/Naloxone.
- 2. Uporova, J., Peacock, A., & Sutherland, R. (2021). Naloxone cascade of care among a sample of people who inject drugs in Canberra, ACT. Drug Trends Bulletin Series. Sydney: National Drug and Alcohol Research Centre, UNSW Sydney.
- 3. Sutherland, et al., (2022). Australian Drug Trends 2022: Key Findings from the National Illicit Drug Reporting System (IDRS) Interviews. Sydney: National Drug and Alcohol Research Centre, UNSW Sydney. DOI: 10.26190/5czp-af24
- 4. NSW Health. Public Drug Warnings (2021). Retrieved January 12, 2023 from https://www.health.nsw.gov.au/aod/public-drug-alerts/Pages/2021.aspx

Suggested citation

Uporova, J., Peacock, A. & Sutherland, R. (2023). Naloxone knowledge among a sample of people that regularly consume illicit stimulants in Australia, 2022. Drug Trends Bulletin Series. Sydney: National Drug and Alcohol Research Centre, UNSW Sydney. DOI: 10.26190/2yyv-3214

Funded by the Australian Government Department of Health and Aged Care under the Drug and Alcohol Program ©NDARC, UNSW SYDNEY 2023. This work is copyright. You may download, display, print and reproduce this material in unaltered form only (retaining this notice) for your personal, non-commercial use or use within your organisation. All other rights are reserved. Requests and enquiries concerning reproduction and rights should be addressed to the information manager, NDARC, UNSW Sydney, NSW 2052, Australia via drugtrends@unsw.edu.au Acknowledgements: The participants who contributed their valuable time to share their experiences, all those who contributed to, advised on, and supported the project and the Australian Government Department of Health and Aged Care for funding the EDRS project.













